

Stochastic stability versus localization in chaotic dynamical systems

Michael Blank ^{*}, Gerhard Keller

Mathematisches Institut, Universitat Erlangen-Nurnberg
Bismarckstrasse 1 1/2, D-91054 Erlangen, Germany.

February 5, 2008

Abstract – We prove stochastic stability of chaotic maps for a general class of Markov random perturbations (including singular ones) satisfying some kind of mixing conditions. One of the consequences of this statement is the proof of Ulam’s conjecture about the approximation of the dynamics of a chaotic system by a finite state Markov chain. Conditions under which the localization phenomenon (i.e. stabilization of singular invariant measures) takes place are also considered. Our main tools are the so called bounded variation approach combined with the ergodic theorem of Ionescu-Tulcea and Marinescu, and a random walk argument that we apply to prove the absence of “traps” under the action of random perturbations.

1 Introduction

The investigation of stochastic stability of chaotic dynamical systems has a long history, and in this paper we shall restrict ourselves mainly to the case of one-dimensional piecewise expanding maps (i.e. Lasota-Yorke [15] type maps). It is well known that if the expanding constant is larger than 2 a Lasota-Yorke type map is stable with respect to mixing random perturbations. It is also known that the assumption that the expanding constant is larger than 1 is not enough for stochastic stability. The point is, only when the expanding constant is larger than 2, the map is really locally expanding, i.e. the image of any small enough interval has Lebesgue measure larger than the genuine one. We prove that for a very general class of Markov random perturbations the mixing condition on perturbations is sufficient for stochastic stability. One of the consequences of this statement is the proof of so called Ulam’s conjecture, about the approximation of a chaotic system by a finite state Markov chain. Conditions under which the localization phenomena (i.e. stabilization of singular invariant measures) takes place are also considered. The nature of this phenomena is a mixing of unstable and stable directions under the action of arbitrary small perturbations. To show that the localization is not the result of discontinuities in the considered class of maps we prove the localization under the action of small mixing random perturbations for general smooth hyperbolic d -dimensional maps.

In this paper we discuss some unusual phenomena that appear in chaotic dynamical systems. First of all, what does it mean a *chaotic* system or map. We shall consider only discrete time *dynamical systems*, i.e. pairs (f, X) , where $X \subset \mathbb{R}^d$ is a compact phase space (say $X = [0, 1]^d$) and $f : X \rightarrow X$ is a nonsingular map, iterations of which define trajectories of the dynamical system. Nonsingular means that $m(f^{-1}A) > 0$ for any measurable set $A \subseteq X$ with positive Lebesgue measure $m(A) > 0$.

^{*}On leave from Russian Academy of Sciences, Inst. for Information Transmission Problems, Ermolovoy Str. 19, 101447, Moscow, Russia, blank@obs-nice.fr

We shall be interested mainly in asymptotic, when time tends to infinity, properties of systems under consideration.

Consider now small *random perturbations* of the discrete time dynamical system. Before giving a rigorous definition of random perturbations of dynamical systems we mention one example, which will be useful to have in mind later on. Let $U \subset R^d$ be a neighborhood of a compact invariant set of the map f such that $fU \subset U$. Suppose that, when we apply f to a point $x \in U$, rather than choosing the exact value of fx we choose in a random way, in accordance with a homogeneous distribution, a point from the ball $B_\varepsilon(fx)$ (i.e. from the ball with centre at the point fx and radius ε). The resulting random map corresponds to independent homogeneously distributed random perturbations.

Definition 1.1 Let $Q_\varepsilon(x, A)$ be a family of transition probabilities and $f : X \rightarrow X$ a map. We denote by f_ε the Markov process on the phase space X , defined by transition the probabilities $Q_\varepsilon(fx, A)$ and call f_ε a random perturbation of f .

We are mainly interested in *small* perturbations, small in the sense that

$$\|Q_\varepsilon h - h\|_1 \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \text{ for any } h \in L^1, \quad (1.1)$$

where $\|\cdot\|_1$ denotes the L^1 -norm. Additionally we assume that our perturbations are *local*, i.e.

$$Q_\varepsilon(x, A) = 0 \text{ for any pair } (x, A), \text{ such that } \text{dist}(x, A) > \varepsilon. \quad (1.2)$$

Therefore the parameter ε here plays the role of a “magnitude” of the perturbation. However, this local condition does not imply stochastic stability even for very smooth chaotic maps as we show in Lemma 2.1. Thus to obtain some kind of stochastic stability the perturbations should satisfy at least some kind of mixing condition (say, for example, of Doeblin’s type [7]). One of the most general assumptions of this type for random perturbations was proposed in [5]:

$$\text{var}(Q_\varepsilon h) \leq \text{var}(h) + C\|h\|_1 \quad (\text{regularity assumption}) \quad (1.3)$$

for any function h of bounded variation ($\text{var}(h) < \infty$). Remark, that for independent random perturbations the transition density depends only on the difference between its arguments: $q_\varepsilon(x, y) = q_\varepsilon(x - y)$, and the constant C on the right hand side of the inequality (1.3) may be set to zero. Thus this condition means that the perturbation is not too far from the independent one. Besides to convolutions with smooth transition probabilities our results apply to many other types of perturbations like bistochastic smooth perturbations, singular perturbations of point mass type and deterministic perturbations by chaotic maps close to identity.

The basic idea of our approach is the following: Consider the Perron-Frobenius operator P , describing the dynamics of densities under the action of the map f , on Banach space $(\text{BV}, \|\cdot\|_{\text{BV}})$ of functions of bounded variation. Here $\|h\|_{\text{BV}} = \text{var}(h) + \|h\|_1$. We prove that the transition operator $P_\varepsilon = Q_\varepsilon P$ of the randomly perturbed map satisfies the uniform Lasota-Yorke type inequality

$$\|P_\varepsilon^N h\|_{\text{BV}} \leq \alpha \cdot \|h\|_{\text{BV}} + C \cdot \|h\|_1 \quad (h \in \text{BV}) \quad (1.4)$$

for some fixed integer N , $\alpha \in (0, 1)$ and $C > 0$ independent of ε . This yields at once the existence of f_ε -invariant densities h_ε with $\|h_\varepsilon\|_{\text{BV}} \leq \frac{C}{1-\alpha}$, such that (1.1) forces each limit $h_* = L^1 - \lim_{\varepsilon \rightarrow 0} h_\varepsilon$ to be an invariant density for f . If the expanding constant λ is larger than 2, (1.4) was proved with $N = 1$ in various settings, see *e.g.* [11, 2, 12] and references therein. For quite a while it was supposed that the extension of this inequality to cases with $\lambda \in (1, 2]$ is only a technical problem. However, the counterexample constructed in [4] shows that the situation is not so simple. After this counterexample (which we shall discuss in Section 2) it became clear, that the main problem is the possible existence of periodic *turning* points, *i.e.* the points, where the derivative of the map f is not well defined. Namely, under the action of random perturbations “traps” or “absorbing sets” can appear near these periodic turning points, which leads to the appearance of new ergodic components in the perturbed

system. In [1] stochastic stability was proved for convolution type smooth random perturbations in situations with $\lambda \in (1, 2]$ but without periodic turning points. However it is not clear whether their argument works for general Markov perturbations.

The main result obtained in these papers is the stability of statistical properties of the perturbed systems in the zero noise limit. We shall show that in the general case there may be generalized phase transitions of type localization - delocalization in such systems. These phase transitions correspond to situations, when trajectories, which should normally be dense, remain confined to a small region (which vanishes, when the coupling constant goes to zero). We call this “localization phenomenon”. The first observation of this type was published in [4].

Definition 1.2 *Let $X = [0, 1]$. A map $f : X \rightarrow X$ is piecewise expanding (PE) if there exists a partition of X into disjoint intervals $\{X_j\}$, such that $f|_{\text{Clos}(X_j)}$ is a C^2 -diffeomorphism (from the closed interval $\text{Clos}(X_j)$ to its image), if the expanding constant of the map*

$$\lambda_f := \inf_{j, x \in X_j} |f'(x)|,$$

is positive, and if $\lambda_{f^\kappa} > 1$ for some iterate f^κ .

For some of our results we impose some extra conditions on f . In particular we distinguish between three different “hyperbolicity” assumptions:

- $f \in H_\infty$, if f is a general PE map;
- $f \in H_1$, if f is a PE map with $\lambda_f > 1$;
- $f \in H_\infty^\varepsilon$ if f is a PE map for which there is $\lambda > 1$ such that $\prod_{k=1}^n |f'_\varepsilon(f_\varepsilon^k x)| \geq \lambda^n$ for each ε -trajectory $(f_\varepsilon^k x)_{k=1,2,\dots}$.

Clearly $H_1 \subset H_\infty^\varepsilon \subset H_\infty$.

Typical examples of PE maps are shown in Figure 1. Observe that maps that we consider need not to be continuous. As it is well known, starting from the paper of Lasota & Yorke [15], these maps have all the statistical properties that one can reasonably expect from deterministic dynamical system. They have a smooth (absolutely continuous) invariant measure μ_f (Sinai-Bowen-Ruelle measure of this map), exponential correlation decay and they obey a CLT with respect to this measure, etc [9, 2]. In this paper we want to discuss stability of these properties with respect to small random perturbations. We shall emphasize some aspects of this problem, because they seem quite counterintuitive, at least from our point of view.

Definition 1.3 *The image of a measure μ under the action of a map f is the measure $f\mu$ defined by $f\mu(A) = \mu(f^{-1}A)$ for any measurable set A . By $f_\varepsilon\mu$ we mean the measure $f_\varepsilon\mu(A) = \int Q_\varepsilon(fx, A)d\mu(x)$. A measure μ is f (f_ε)-invariant, if $f\mu = \mu$ ($f_\varepsilon\mu = \mu$). μ is called smooth, if it has a density with respect to Lebesgue measure.*

Definition 1.4 *A probability measure μ_f is called a Sinai-Bowen-Ruelle (SBR) measure of f , if there exists an open subset U of the phase space such that the images $f^n\mu$ of any smooth probability measure μ with the support in U converge weakly to μ_f . Analogously we define SBR measures μ_ε for perturbed systems f_ε .*

Definition 1.5 *A turning point of a map f is a point, where the derivative of the map is not well defined. The set of turning points is denoted by TP, the set of periodic turning points by PTP. (In Figure 1a we have $c \in \text{TP}$, in Figure 1b we have $c \in \text{PTP}$.)*

Standing assumption: From now on we assume that all perturbations Q_ε satisfy (1.1), (1.2) and (1.3).




Figure 1: PE maps. (a) general PE map, (b) W-map

Theorem 1.1 [4, 11] *Suppose $\lambda_f > 2$ and that f has a unique smooth SBR measure μ_f . Then for any sufficiently small $\varepsilon > 0$ also the perturbed system f_ε has a unique smooth invariant SBR measure μ_ε , and the μ_ε converge weakly to μ_f as $\varepsilon \rightarrow 0$.*

In this paper we generalize Theorem 1.1 and show that at least some additional assumptions are necessary for the stability of the smooth SBR measure μ_f .

Theorem 1.2 *Let $f \in H_\infty$, $\text{PTP} = \emptyset$, and suppose that f has a unique smooth SBR measure μ_f . Then for any $\varepsilon > 0$ small enough the perturbed system f_ε has a smooth invariant SBR measure μ_ε converging weakly as $\varepsilon \rightarrow 0$ to the smooth SBR measure μ_f .*

So the only topological obstacle for the stability of statistical properties of PE maps under small perturbations is the existence of periodic turning points. Consequently, generic PE maps are stochastically stable. However, in order to investigate stability properties of families of maps, rather than individual maps, also maps with periodic turning points must be studied.

Theorem 1.3 *There exists a map $f \in H_1$ with periodic turning points and smooth perturbations Q_ε such that for sufficiently small $\varepsilon > 0$ the perturbed system f_ε has a unique invariant SBR measure μ_ε , converging to a singular and unstable invariant measure of the map f as $\varepsilon \rightarrow 0$.*

This theorem does not only state the localization of f_ε -invariant measures (supports on small sets), but also guarantees their smoothness and the absence of other SBR measures. Actually the localization phenomenon was firstly shown in [4, 5], but the statement about the smoothness of invariant measures, absence of other SBR measures, and investigation of their properties are new.

The nature of the localization phenomenon here is due to the fact that if the expanding constant λ_f is less than 2 the map f is not locally expanding near periodic singular points. Indeed it may map a small interval Δ to the interval of length $\lambda_f|\Delta|/2$. This property distinguishes the situations in Theorems 1.1 and 1.2. To show that this phenomenon is not something obscure, specific for only discontinuous maps, we shall prove its presence for a general multidimensional smooth hyperbolic map in Section 2.

In order to exclude the behaviour described in Theorem 1.3 we introduce the following *random walk (RW)* assumption:

$$\text{A family } Q_\varepsilon \text{ } (\varepsilon > 0) \text{ belongs to the class } RW, \text{ if there are } 0 < \theta, \delta < 1 \text{ such that} \quad (1.5)$$

$$Q(x, [0, x - \theta\varepsilon]) > \delta \text{ and } Q(x, [x + \theta\varepsilon, 1]) > \delta \text{ for all } x.$$

For the next two theorems we assume additionally that a local version of (1.3) is satisfied, see (3.6).

Theorem 1.4 *Let $f \in H_\infty$, PTP not necessarily empty, and suppose that the transition probabilities Q_ε belong to RW and have densities $q_\varepsilon(x, y)$ satisfying*

$$\text{var}(y \mapsto q_\varepsilon(x, y)) \leq \frac{M}{\varepsilon} \quad \text{for all } x.$$

Then for any sufficiently small $\varepsilon > 0$ the perturbed system f_ε has a smooth invariant SBR measure μ_ε converging weakly to the smooth SBR measure μ_f as $\varepsilon \rightarrow 0$.

To formulate results about more general singular perturbations we need the notion of renormalizability, which will be discussed in detail at the end of Section 3.3. Roughly speaking this notion means that for any $\varepsilon > 0$ small enough a map (or a perturbation) can be renormalized to a fixed shape in small neighbourhoods (of order ε) of periodic turning points.

Theorem 1.5 *Let $f \in H_\infty^\varepsilon$, PTP not necessarily empty, and $Q \in \text{RW}$. Then for any $\varepsilon > 0$ small enough the perturbed system f_ε has a smooth invariant SBR measure μ_ε and the transition operator $Q_\varepsilon P_f$ satisfies the Lasota-Yorke type inequality. If additionally both the map and the perturbations are renormalizable, then the measures μ_ε converge weakly as $\varepsilon \rightarrow 0$ to the smooth SBR measure μ_f .*

The last result of the paper is the proof of Ulam's conjecture [18] on the approximation of chaotic dynamics by finite state Markov chains. The idea of the construction is to take a finite partition $\{\Delta_i\}$ of the phase space with bounded volume ratios and to approximate the action of the map f by the Markov chain with transition probabilities

$$p_{ij} := \frac{|\Delta_i \cap f^{-1}\Delta_j|}{|\Delta_i|}.$$

This construction could be considered as a special type of small random perturbations. In fact, our standing assumption on Q_ε is satisfied for this particular class of perturbations. The correctness of Ulam's conjecture was proved in [14] for PE maps with $\lambda_f > 2$. In contrast to our other statements the answer to the conjecture does not depend on the existence of periodic turning points.

Theorem 1.6 *Let $f \in H_\infty$. Then invariant measures constructed by Ulam's procedure converge weakly to the smooth SBR measure μ_f .*

2 Counterexamples

2.1 Localization in the absence of the regularity assumption

Our first example shows that even for arbitrarily smooth chaotic maps and absolutely continuous perturbations the *local* condition (1.2) alone does not imply stochastic stability.

Lemma 2.1 *Suppose $f : X \rightarrow X$ has a cycle (periodic trajectory) $\bar{c} = c_1, \dots, c_n$ and satisfies a Lipschitz condition in some neighborhood of this cycle. Then there exists a family of local random perturbations with densities $q_\varepsilon(x, y)$ for which the f_ε have smooth invariant measures μ_ε converging to the δ -measure on this cycle when $\varepsilon \rightarrow 0$.*

Proof. Fix a constant β greater than the local Lipschitz constant of the map f and consider a family of transition probability densities

$$q_\varepsilon(x, y) = \begin{cases} 1/|B_{\varepsilon/\beta}(\bar{c})|, & \text{if } x \in B_\varepsilon(\bar{c}) \text{ and } y \in B_{\varepsilon/\beta}(\bar{c}); \\ 0, & \text{if } x \in B_\varepsilon(\bar{c}) \text{ and } y \notin B_{\varepsilon/\beta}(\bar{c}); \\ 1/|B_\varepsilon(\bar{c})|, & \text{if } x \notin B_\varepsilon(\bar{c}) \text{ and } y \in B_\varepsilon(\bar{c}); \\ 0, & \text{otherwise.} \end{cases}$$

Then clearly f_ε has a new ergodic component in the ε -neighborhood of the cycle $\bar{c}$, because $fB_{\varepsilon/\beta}(\bar{c}) \subset B_\varepsilon(\bar{c})$ and by the construction $q_\varepsilon(x, y) = 0$ for $x \in B_\varepsilon(\bar{c})$ and $y \notin B_{\varepsilon/\beta}(\bar{c})$. $\blacksquare$

This proof shows that for stochastic stability some additional assumption is needed that prevents the perturbation to act precisely against the dynamics of the map. This is one of the purposes of the regularity assumption (1.3).

2.2 Periodic turning points and the RW assumption, proof of Theorem 1.3.

Fix a number $10 < \beta < \infty$ and consider the following family of transition probability densities:

$$q_\varepsilon(x, y) = \begin{cases} \frac{\beta}{2\varepsilon}, & \text{if } (1 - \frac{1}{\beta})\varepsilon \leq -x + y \leq (1 + \frac{1}{\beta})\varepsilon; \\ 0, & \text{otherwise.} \end{cases} \quad (2.1)$$

Note that random perturbations with such transition densities are of convolution type and have a very strong mixing property - exponential rate of correlation decay. The following lemma proves at the same time Theorem 1.3:

Lemma 2.2 ([5]) *Let $1 < \lambda < 2 - 1/\beta$, and let the map f is defined as follows:*

$$f(x) = \begin{cases} 1 - \frac{2x}{1-1/\lambda}, & \text{if } 0 \leq x \leq (1 - 1/\lambda)/2; \\ \lambda x - \frac{\lambda-1}{2}, & \text{if } (1 - 1/\lambda)/2 < x \leq 1/2; \\ f(1-x), & \text{otherwise.} \end{cases} \quad (2.2)$$

See Figure 1.b. Then for all sufficiently small $\varepsilon > 0$ the perturbed map f_ε has a unique invariant measure converging to the δ -measure at $\frac{1}{2}$ as $\varepsilon \rightarrow 0$.

The **proof** of this statement is based on the fact that trajectories of f_ε starting from $[1/2 - \varepsilon(\lambda(\beta + 1) - \beta + 1)/\beta, 1/2 + \varepsilon(\beta + 1)/\beta]$ remain in this interval for ever with probability 1. A related example with the same map f but different transition kernel was given in [11].

We remark that, as $|f'(x)| = \lambda < 2$ in some neighborhood of the fixed point $c \in X$, the mapping f is not expanding in some sense, because it maps an interval neighborhood U of the fixed point to the interval fU where

$$|fU| \leq \frac{\lambda|U|}{2} < |U|.$$

So, in the presence of periodic turning points, instability of these points under f_ε is a necessary extra requirement. For this reason we introduced the random walk assumption RW .

In a broader context the inequality $|fU| < |U|$ can be interpreted as a mixing of stable and unstable directions under the action of random perturbations. In fact, a more general statement is:

Theorem 2.1 *Let a C^2 -smooth map $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$ has a hyperbolic fixed point $c \in \mathbb{R}^d$, such that for the linear part $D_c f$ at this point we have*

$$\mathbb{R}^d = E_c^s + E_c^u, \quad \dim(E_c^s) \geq \dim(E_c^u) = N,$$

$$|D_c f(x)| \leq \lambda_s |x| \quad \text{for all } x \in E_c^s,$$

$$\lambda_u |x| \leq |D_c f(x)| \leq \Lambda_u |x| \quad \text{for all } x \in E_c^u,$$

and $\lambda_s \Lambda_u < 1$. Then there exist a family of local mixing stochastic ε -perturbations such that for all $\varepsilon > 0$ small enough the stochastically ε -perturbed map f_ε has a smooth invariant measure in small neighborhood of the point c , converging to the delta-measure at this point as $\varepsilon \rightarrow 0$.

Proof. Let $\tau_\alpha : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a d -dimensional rotation around the point $c \in \mathbb{R}^d$ through the angle α , such that $\tau_\alpha E_c^u \subset E_c^s$ and any coordinate of the vector α lies in the region $[-\pi/2, \pi/2]$. For each $\varepsilon > 0$ we define a map $\tau_\alpha^{(\varepsilon)} : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that for all $x \in B_{\varepsilon/\sqrt{N}}(c)$, where $N = \dim(E_c^u)$, it coincides with τ_α and in the complement of this ball it is defined as follows:

$$\tau_\alpha^{(\varepsilon)} x = \tau_{\varepsilon\alpha/(|x-c|\sqrt{N})} x.$$

Then

$$|x - \tau_\alpha^{(\varepsilon)} x| \leq \varepsilon\pi/2.$$

Now we fix some $0 < \beta \ll 1$ and define the following family of transition probability densities:

$$q_\varepsilon(x, y) = \begin{cases} 1/|B_{\beta\varepsilon}|, & \text{if } y \in B_{\beta\varepsilon}(\tau_\alpha^{(\varepsilon)} x); \\ 0, & \text{otherwise.} \end{cases}$$

Evidently, that such perturbations are mixing and due to the estimate above $q_\varepsilon(x, y) = 0$ for $|x - y| > 2\varepsilon$ and $\beta < 0.4$. On the other hand just as in Lemma 2.2 any trajectory of the stochastically ε -perturbed map, started from the ball $A_\varepsilon = \{x \in \mathbb{R}^d : |x - c| \leq \varepsilon/(2\Lambda_u)\}$ remains in it with the probability 1. Indeed for

$$\beta < 2\Lambda_u(1 + \lambda_s\Lambda_u)/(1 - \lambda_s\Lambda_u)$$

we have

$$\lambda_s\Lambda_u(\varepsilon/(2\Lambda_u) + \beta\varepsilon) + \Lambda_u\beta\varepsilon < \varepsilon/(2\Lambda_u),$$

which finishes the proof. Note that the constant β here does not depend on ε . ■

The generalization of this result for an arbitrary hyperbolic periodic trajectory is straightforward.

2.3 H_∞ versus H_∞^ε assumption

In the case of singular perturbations we have to replace the simple hyperbolicity assumption H_∞ by the stronger one H_∞^ε , see Theorems 1.4 and 1.5. To construct an example satisfying H_∞ without being stochastically stable, consider a map f with fixed turning point $c = f(c)$, such that locally (in a neighborhood of this point) the map is defined as follows:

$$f(x) := \begin{cases} \lambda_1 x + (1 - \lambda_1)c, & \text{if } x \leq c \\ \lambda_2 x + (1 - \lambda_2)c, & \text{otherwise,} \end{cases} \quad (2.3)$$

where $|\lambda_1| > 1 > |\lambda_2| > 0$ and $|\lambda_1\lambda_2| > 1$. Both of the maps in Figure 2 satisfy these assumptions. We consider the following singular perturbations:

$$x \longrightarrow \begin{cases} x + \varepsilon, & \text{with probability } 1 - q \\ x - \varepsilon, & \text{with probability } q, \end{cases} \quad (2.4)$$

where $0 < q < 1$ is a parameter. Note that this setting satisfies the RW assumption.

Clearly, the map $f(x) + \varepsilon$ has a stable fixed point $c_\varepsilon := c + \varepsilon/(1 - \lambda_2)$. Now let $h \in L^1$ be a nonnegative function, such that $h(c_\varepsilon) > 0$. Then the transition operator P of the random map satisfies the following inequality:

$$Ph(c_\varepsilon) \geq (1 - q) \left(\frac{1}{|\lambda_1|} h\left(c + \frac{\varepsilon}{1 - \lambda_1}\right) + \frac{1}{|\lambda_2|} h(c_\varepsilon) \right) \geq \frac{1 - q}{|\lambda_2|} h(c_\varepsilon).$$

Therefore for any positive integer n we have

$$P^n h(c_\varepsilon) \geq \left(\frac{1 - q}{|\lambda_2|} \right)^n h(c_\varepsilon).$$




Figure 2: PE maps. (a) nonsymmetric W-map, example of a PE map of type H_∞

Thus, for $1 - q > |\lambda_2|$ the value of $P^n h$ at the point c_ε is growing exponentially, so an exponential decay assumption in L^∞ - or BV-norm that we introduce in Section 3.3.1 is violated.

In spite of this, using different methods, we are able to prove the convergence to a smooth SBR measure μ_ε in this example. Note, that new phenomenon is observed here. Indeed, due to the estimates above, the density of the SBR measure μ_ε is not a function of bounded variation and usual Lasota-Yorke type estimates do not work here. The point is that the mathematical expectation of the slope of the random map at some points is less than 1 (in our example $E(|f'(c_\varepsilon)|) = |\lambda_2| < 1$), which contradicts usual settings (see [16, 17] and references therein). Indeed, if such a point is a fixed point for some of the shifted maps (as the point c_ε for the map $f(x) + \varepsilon$ in our example), then this really leads to unusual properties of the transition operator of the random map. Investigation of ergodic properties of such random maps is out of the scope of the present paper and will be published elsewhere.

3 Proofs of the stability theorems

3.1 The general setting

3.1.1 The deterministic part of the dynamics

Let X be a bounded interval in $\mathbb{R}$, and denote by m Lebesgue measure on X . Three function spaces over X play an important role in our studies:

- $L^1 = L_m^1(X)$, the space of m -equivalence classes of complex valued integrable functions on X , which is endowed with the norm $\|h\|_1 = \int_X |h| dm$,
- $L^\infty = L_m^\infty(X)$, the space of m -equivalence classes of complex valued bounded functions on X , which is endowed with the norm $\|h\|_\infty = \text{ess sup}_X |h|$, and
- BV, the space of m -equivalence classes of integrable functions of bounded variation on X . It is endowed with the norm $\|h\|_{\text{BV}} = \|h\|_1 + \text{var}(h)$ where $\text{var}(h) = \text{var}_{\mathbb{R}}(h)$ and where for any $Y \subseteq \mathbb{R}$

$$\begin{aligned} \text{var}_Y(h) &= \sup \left\{ \int_X \varphi' h dm : \varphi \in \mathcal{T}_1(Y) \right\} \quad \text{and} \\ \mathcal{T}_1(Y) &= \{ \varphi \in C(\mathbb{R}) : \|\varphi\|_\infty \leq 1, \varphi|_{\mathbb{R} \setminus Y} = 0, \varphi \text{ differentiable, } \|\varphi'\|_\infty < \infty \} . \end{aligned}$$

We remark that the same notion of variation is obtained if one restricts $\mathcal{T}_1(Y)$ to C^1 - or even C^∞ -functions.

Remark 3.1 As X is bounded, $\|h\|_\infty \leq \frac{1}{2} \text{var}(h)$ for all $h \in \text{BV}$, and if $I \subseteq X$ is an interval, then $\text{var}(h \cdot 1_I) \leq \text{var}(h)$. Indeed, our setting means that $\text{var}(h)$ is the variation of $1_X \cdot h$ over $\mathbb{R}$.

We study dynamics on X given by a map $f : X \rightarrow X$ composed with a random perturbation with transition kernel Q . More specifically, let $f : X \rightarrow X$ be piecewise monotone with a finite number of turning points $\text{TP} := \{c_1, \dots, c_r\}$. By this we mean that for each maximal component Z of $X \setminus \text{TP}$ holds

$$\begin{aligned} f|_Z \text{ is monotone and continuously differentiable and extends continuously to } \bar{Z}, \\ \text{var}_Z |f'| < \infty. \text{ In particular, } \Lambda := \sup_X |f'| < \infty. \end{aligned} \quad (3.1)$$

These intervals Z form a partition $\mathcal{Z}$ of X modulo the finite set TP . Let $\mathcal{Z}_n = \{Z_1 \cap f^{-1}Z_2 \cap \dots \cap f^{-(n-1)}Z_n \neq \emptyset : Z_i \in \mathcal{Z}\}$. Then $\mathcal{Z}_n$ is a partition of X modulo finitely many points, and $f|_Z^n$ satisfies the basic assumptions (3.1) for each $Z \in \mathcal{Z}_n$.

Our essential hyperbolicity assumption is:

There are constants $\lambda > 1$, $\eta > 0$ and $N \in \mathbb{N}$ such that

$$|(f|_Z^N)'| \geq \lambda \text{ for all } Z \in \mathcal{Z}_N \quad \text{and} \quad \eta \leq |f'| \leq \eta^{-1} \text{ for all } Z \in \mathcal{Z}. \quad (3.2)$$

Of course, once we have such a constant $\lambda > 1$, we can (and will) assume that (for a larger iterate N) we even have $\lambda > 4$.

Given such an iterate N and a suitable $\beta > 0$ that we specify later, we refine the partition $\mathcal{Z}$ by adding further points to TP in such a way that

$$\text{var} \left(\frac{f'|_Z}{\eta} \right), \quad \text{var} \left(\frac{\eta^{-1}}{f'|_Z} \right) \leq \beta. \quad (3.3)$$

We further modify the set TP by doubling all its elements together with their preimages and extend f to the enlarged space by one-sided continuity. We call the resulting space, the extended map and the collection of doubled turning points again X , f , and TP respectively. (Strictly speaking the new space X is a linearly ordered, order complete space. Its order topology will be quite different from the topology on the real line, usually it will be totally disconnected.)

If after these modifications there are $c_i, c_j \in \text{TP}$ such that $f^k c_i = c_j$ for some $k > 0$, we also add all points $f^l c_i$, $l = 1, \dots, k-1$, to TP . In this way we can make sure that

$$\text{for each } c_i \in \text{TP} \text{ either } f c_i \in \text{TP} \text{ or } f^k c_i \notin \text{TP} \text{ for all } k \geq 1. \quad (3.4)$$

We denote the enlarged set TP once more by $\text{TP} = \{c_1, \dots, c_r\}$. For $c_j \in \text{TP}$ we mean by $c_j \in \text{TP}$ its doubled copy.

Let $J_j := [c_j, d_j] \subset \tilde{J}_j := [c_j, \tilde{d}_j]$, $j = 1, \dots, r$ be two families of one-sided interval neighbourhoods of the turning points. Here as in the sequel we write $[u, v] = \{x \in X : u \leq x \leq v \text{ or } v \leq x \leq u\}$, in particular $[u, v] = [v, u]$. The $\tilde{J}_j$ are chosen such that any two of them are disjoint. (Observe that this holds also for $\tilde{J}_j$ and $\tilde{J}_j$!) We write $Y = \cup_j J_j$ and $\tilde{Y} = \cup_j \tilde{J}_j$ and assume that for each $j = 1, \dots, r$ holds

- either $f(J_j) \cap \tilde{Y} = \emptyset$
 - or $f(J_j) \supset \tilde{J}_i$ for some i in such a way that $f(c_j) = c_i$ but $f(d_j) \notin \tilde{J}_i$.
- (3.5)

Observe that this includes a kind of topological expansion assumption for neighbourhoods of periodic turning points. If $f \in H_\infty$, then this assumption can always be satisfied.

On the level of “mass transport” the dynamics of f are described by the *Perron-Frobenius operator* $P_f : L^1 \rightarrow L^1$,

$$P_f h = \sum_{Z \in \mathcal{Z}} \frac{h}{|f'|} \circ f_{|Z}^{-1} \cdot 1_{fZ} .$$

We note that $\int P_f h \, dm = \int h \, dm$ for all $h \in L^1$. Later we shall see that P_f is also a bounded linear operator on BV.

3.1.2 The stochastic part of the dynamics

Perturbations are described by (sub)-Markovian operators $Q : L^1 \rightarrow L^1$. More precisely we assume

1. Q is a linear, positive operator and $\|Q\|_1 \leq 1$.
2. $Q^* : L^\infty \rightarrow L^\infty$ denotes the dual operator to Q .

We remark that Q can always be thought of as represented by a (sub)-Markov transition kernel, i.e.

$$Q^* f(x) = \int f(y) Q(x, dy) \quad \text{and} \quad Qh(y) = \left(\frac{d}{dm} \int h(x) Q(x, \cdot) m(dx) \right) (y) ,$$

see *e.g.* [13, §3.1]. If $Q(x, \cdot) \ll m$ for each x (what our assumptions do not necessarily imply), we denote $q(x, y) = \frac{d}{dm} Q(x, \cdot)(y)$.

The following regularity assumption on Q plays an essential role: There is a constant $C > 0$ such that

$$\begin{aligned} \text{var}(Qh) &\leq \text{var}(h) + C \cdot \|h\|_1 \quad \text{and} \\ \text{var}_{J_k \cup \tilde{J}_k}(Qh) &\leq \text{var}_{\tilde{J}_k \cup \tilde{J}_k}(h) + C \cdot \|h \cdot 1_{\tilde{J}_k \cup \tilde{J}_k}\|_1 \end{aligned} \tag{3.6}$$

for each $h \in \text{BV}$ and $k \in \{1, \dots, r\}$. As we shall see later, this assumption is satisfied in many cases of interest, including absolutely continuous, random walk-like and also deterministic perturbations.

Finally we define the *spread* of Q as

$$\text{spread}(Q) = \inf\{\delta > 0 : Q(x, \{y : |y - x| > \delta\}) = 0 \, \forall x\} .$$

The spread is the size of the largest jump that can be caused by the perturbation. We assume that

$$\text{spread}(Q) < \min\{|\tilde{d}_j - d_j| : j = 1, \dots, r\} , \tag{3.7}$$

i.e. random jumps from $X \setminus \tilde{Y}$ to Y are excluded.

3.1.3 The decomposition

In the sequel we study the behaviour of f on Y and on $X \setminus Y$ separately under the assumption that transitions from $X \setminus Y$ to Y are restricted in the following sense: Define $\tilde{P}_1, \tilde{P}_2 : BV \rightarrow BV$ by

$$\tilde{P}_1 h = QP_f(h \cdot 1_{X \setminus Y}), \quad \tilde{P}_2 h = QP_f(h \cdot 1_Y) . \tag{3.8}$$

A straightforward calculation shows

Proposition 3.1 *Suppose that there are constants $C_1, C_2 > 0$ and $\alpha \in (0, 1)$ such that*

$$\text{var}(\tilde{P}_j^k h) \leq C_1 \alpha^k \text{var}(h) + C_2 \|h\|_1 \quad \text{for all } k \in \mathbb{N} \tag{3.9}$$

and that there is some $N \in \mathbb{N}$ such that

$$\tilde{P}_2 \tilde{P}_1^k \tilde{P}_2 = 0 \quad \text{for all } k = 1, \dots, N. \quad (3.10)$$

Then

$$\text{var}((QP_f)^N h) \leq \left(\frac{N(N+1)}{2} C_1^3 + C_1 \right) \alpha^N \cdot \text{var}(h) + (1 + C_1 + C_1^2) C_2 \frac{(N+1)^2}{2} \cdot \|h\|_1. \quad (3.11)$$

In order to guarantee (3.10) for a fixed N the intervals J_j and the spread of Q must be taken sufficiently small:

Lemma 3.2 *Given f , N and the set TP , there is $\delta > 0$ such that (3.10) is satisfied if $\text{spread}(Q) < \delta$ and $|J_j| < \delta$ for all $j = 1, \dots, r$.*

Proof. Consider a trajectory of the perturbed system $x_0, x_1, \dots, x_k$ that starts at $x_0 \in J_i \subseteq Y$ and suppose that it ends at $x_k \in J_j \subseteq Y$ with $x_l \notin X \setminus Y$ for $l = 1, \dots, k-1$. If δ in the assumptions of the lemma is small, the trajectory stays close to a sequence $c_i = z_0, z_1, \dots, z_k$ satisfying for each $l = 1, \dots, k$ either $z_l = f z_{l-1}$ or $z_l \in \{c_m, c_{\tilde{m}}\}$ if $c_m = f z_{l-1}$. In particular, making δ small we can force x_k to be as close as we like to the point $z_k \in \cup_{l=1}^k f^l(\text{TP})$, and as $x_k \in J_i$, z_k is at the same time close to c_j . Because of (3.4) this is possible only if $z_0, \dots, z_k \in \text{TP}$. But this is incompatible with the assumptions (3.5) and (3.7). $\blacksquare$

3.1.4 The scheme of the proofs

Given not only one perturbation operator but a whole family Q_ε ($\varepsilon > 0$), we will have to show that there are decompositions $X = Y \cup (X \setminus Y)$ of X depending on ε , such that the corresponding operators $\tilde{P}_{1,\varepsilon}$ and $\tilde{P}_{2,\varepsilon}$ satisfy (3.9) uniformly in ε (i.e. with constants C_1, C_2, α not depending on ε). Then there is $N > 0$ such that $\left(\frac{N(N+1)}{2} C_1^3 + C_1 \right) \alpha^N < \frac{1}{2}$, and according to Lemma 3.2 and Proposition 3.1 there are constants $\delta, C_3 > 0$ such that

$$\text{var}((Q_\varepsilon P_f)^N h) \leq \frac{1}{2} \cdot \text{var}(h) + C_3 \cdot \|h\|_1 \quad \text{for all } h \in \text{BV}. \quad (3.12)$$

As usual this implies that each $Q_\varepsilon P_f$ has an invariant density h_ε and $\text{var}(h_\varepsilon) \leq 2C_3$ for all ε . In particular, the family $(h_\varepsilon)_{\varepsilon > \delta}$ is relatively compact in L^1 . Consider any limit $h = \lim_{i \rightarrow \infty} h_{\varepsilon_i}$ with $\varepsilon_i \rightarrow 0$. The invariance $Q_\varepsilon P_f h_\varepsilon = h_\varepsilon$ follows immediately from

$$\|P_f h - h\|_1 \leq \|P_f h - E_\varepsilon P_f h\|_1 + \|Q_\varepsilon P_f h - Q_\varepsilon P_f h_\varepsilon\|_1 + \|h_\varepsilon - h\|_1$$

and the assumption that $\|Q_\varepsilon P_f h - P_f h\|_1 \rightarrow 0$ as $\varepsilon \rightarrow 0$. We specialize this to

Lemma 3.3 *If P_f has a unique invariant density h , then, under the above assumptions on the family Q_ε , the invariant densities h_ε of $Q_\varepsilon P_f$ converge to h in L^1 as $\varepsilon \rightarrow 0$.*

This lemma reduces the proofs of Theorems 1.2, 1.4, 1.5 and 1.6 to proving (3.9) uniformly in ε under the various assumptions of these four theorems. Much more detailed informations about the convergence $h_\varepsilon \rightarrow h$ and about the ergodic properties of the perturbed systems f_ε can be gained by exploiting more carefully the spectral theoretic consequences of (3.12) in the light of the Ionescu-Tulcea/Marinescu theorem, see e.g. [11].

3.2 Estimates far from turning points

In this section we prove (3.9) for $\tilde{P}_1$, i.e. for branches with orbits far from turning points. As an immediate corollary we obtain Theorem 1.2.

3.2.1 Interchanging the map and the perturbation

Fix a sequence of monotone branches $f_i = f|_{Z_i \setminus Y}$, $i = 1, \dots, N$ of f and study the operator $QP_{f_1}QP_{f_2}\dots QP_{f_N}$. For each of the $i = 1, \dots, N$ we fix a monotone and continuously differentiable extension $\tilde{f}_i : \mathbb{R} \rightarrow \mathbb{R}$ of f_i . Given $\beta > 0$ as in (3.3) one can always find such an extension with the additional property that there is $y_0 \in Z_i$ such that

$$\text{var}\left(\frac{\tilde{f}_i'}{f'(y_0)}\right), \text{var}\left(\frac{f'(y_0)}{\tilde{f}_i'}\right) \leq \beta. \quad (3.13)$$

Lemma 3.4 *Let Q be a (sub)Markov operator satisfying*

$$\text{var}(Qh) \leq \text{var}(h) + C \cdot \|h\|_1 \text{ for all } h \in \text{BV}.$$

For a fixed branch f_i of f as above define $\tilde{Q} := P_{\tilde{f}_i^{-1}}QP_{f_i}$. Then $P_{\tilde{f}_i}\tilde{Q} = QP_{f_i}$ and

$$\text{var}(\tilde{Q}h) \leq (1 + \frac{3}{2}\beta)^2 \cdot \text{var}(h \cdot 1_I) + C(1 + \frac{3}{2}\beta)\|f'\|_\infty \cdot \|h \cdot 1_I\|_1$$

for all $h \in \text{BV}$.

Proof. For notational convenience we write f and $\tilde{f}$ instead of f_i and $\tilde{f}_i$, and we denote $I = Z_i \setminus Y$. Let $h \in \text{BV}$, $y \in X$ and $u(y) := \frac{\tilde{f}'(y)}{f'(y_0)}$. Then $\text{var}(u) \leq \beta$, $\|u\|_\infty \leq 1 + \beta$ and

$$\tilde{Q}h(y) = \frac{1}{|(\tilde{f}^{-1})'(\tilde{f}y)|} \cdot QP_f h(\tilde{f}y) = u(y) \cdot QP_f(f'(y_0) \cdot h)(\tilde{f}y)$$

because QP_f is a linear operator. Therefore

$$\begin{aligned} & \text{var}(\tilde{Q}h) \\ & \leq \text{var}(u) \cdot \|QP_f(f'(y_0) \cdot h)\|_\infty + \|u\|_\infty \cdot \text{var}(QP_f(f'(y_0) \cdot h)) \\ & \leq \beta \cdot \frac{1}{2} \text{var}(QP_f(f'(y_0) \cdot h)) + (1 + \beta) \cdot \text{var}(QP_f(f'(y_0) \cdot h)) \\ & \leq (1 + \frac{3}{2}\beta) \cdot (\text{var}(P_f(f'(y_0) \cdot h)) + C \cdot \|P_f(f'(y_0) \cdot h)\|_1) \\ & \leq (1 + \frac{3}{2}\beta) \cdot \left(\text{var}\left(\left(\frac{f'(y_0)}{\tilde{f}_i'}\right) \cdot h\right) \circ f^{-1} \cdot 1_{fI} + C \cdot \|f'(y_0) \cdot h \cdot 1_I\|_1 \right) \\ & \leq (1 + \frac{3}{2}\beta) \cdot \left(\text{var}\left(\frac{f'(y_0)}{\tilde{f}_i'} \cdot h \cdot 1_I\right) + C \cdot \|f'\|_\infty \cdot \|h \cdot 1_I\|_1 \right) \\ & \leq (1 + \frac{3}{2}\beta) \cdot \left(\text{var}\left(\frac{f'(y_0)}{\tilde{f}_i'}\right) \|h \cdot 1_I\|_\infty + \left\|\frac{f'(y_0)}{\tilde{f}_i'}\right\|_\infty \text{var}(h \cdot 1_I) \right. \\ & \quad \left. + C \cdot \|f'\|_\infty \cdot \|h \cdot 1_I\|_1 \right) \\ & \leq (1 + \frac{3}{2}\beta) \cdot \left(\beta \cdot \frac{1}{2} \text{var}(h \cdot 1_I) + (1 + \beta) \cdot \text{var}(h \cdot 1_I) + C \cdot \|f'\|_\infty \cdot \|h \cdot 1_I\|_1 \right) \\ & \leq (1 + \frac{3}{2}\beta)^2 \text{var}(h \cdot 1_I) + (1 + \frac{3}{2}\beta)C\|f'\|_\infty \cdot \|h \cdot 1_I\|_1 \end{aligned}$$

■

Next we apply this lemma to the sequence $f_1, \dots, f_N$ of branches of f .

Lemma 3.5 Let $Q_1, \dots, Q_N$ be (sub)Markov operators with

$$\text{var}(Q_j h) \leq \text{var}(h) + C_j \|h\|_1$$

for some constants $C_j > 0$ ($j = 1, \dots, N$). Then there is for each $k = 1, \dots, N$ a (sub)Markov operator $\tilde{Q}_k$ such that

$$Q_1 P_{f_1} Q_2 P_{f_2} \dots Q_k P_{f_k} = P_{\tilde{f}_1} P_{\tilde{f}_2} \dots P_{\tilde{f}_k} \tilde{Q}_k$$

and for each $h \in \text{BV}$

$$\text{var}(\tilde{Q}_k h) \leq (1 + \frac{3}{2}\beta)^{2k} \cdot \text{var}(h) + \tilde{C}_k \cdot \|h\|_1$$

where

$$\tilde{C}_k = \sum_{j=1}^k (1 + \frac{3}{2}\beta)^{2(j-1)} C_j \prod_{i=j}^k [(1 + \frac{3}{2}\beta) \|f'_i\|_\infty] .$$

Proof. The proof is by induction. For $k = 1$ this is just the statement of the previous lemma. Suppose the lemma is true for $k = n$. Then

$$Q_1 P_{f_1} Q_2 P_{f_2} \dots Q_n P_{f_n} Q_{n+1} P_{f_{n+1}} = P_{\tilde{f}_1} P_{\tilde{f}_2} \dots P_{\tilde{f}_n} \tilde{Q}_n Q_{n+1} P_{f_{n+1}}$$

by inductive hypothesis. The previous lemma applied to $P_f = P_{f_{n+1}}$ and $Q = \tilde{Q}_n Q_{n+1}$ yields the existence of $\tilde{Q}_{n+1}$ such that $\tilde{Q}_n Q_{n+1} P_{f_{n+1}} = P_{\tilde{f}_{n+1}} \tilde{Q}_{n+1}$. Since

$$\begin{aligned} \text{var}(\tilde{Q}_n Q_{n+1} h) &\leq (1 + \frac{3}{2}\beta)^{2n} \cdot \text{var}(Q_{n+1} h) + \tilde{C}_n \cdot \|Q_{n+1} h\|_1 \\ &\leq (1 + \frac{3}{2}\beta)^{2n} \cdot \text{var}(h) + ((1 + \frac{3}{2}\beta)^{2n} C_{n+1} + \tilde{C}_n) \cdot \|h\|_1 , \end{aligned}$$

we have

$$\text{var}(\tilde{Q}_{n+1} h) \leq (1 + \frac{3}{2}\beta)^{2(n+1)} \text{var}(h) + (1 + \frac{3}{2}\beta) \|f'_{n+1}\|_\infty (\tilde{C}_n + (1 + \frac{3}{2}\beta)^{2n} C_{n+1}) \cdot \|h\|_1 .$$

Inserting the formula for $\tilde{C}_n$ finishes the proof. ■

Now we combine this lemma with the hyperbolicity assumption (3.2) for f :

Lemma 3.6 Suppose that in the situation of the previous lemma there are constants $\lambda > 1$, $\eta > 0$ such that

$$|(f|_Z^N)'| \geq \lambda \text{ for all } Z \in \mathcal{Z}_n \quad \text{and} \quad |f|_Z' \geq \eta \text{ for all } Z \in \mathcal{Z} .$$

(This is assumption (3.2).) If $f_1 \circ \dots \circ f_N$ is not the empty map (i.e. if $f_1 \circ \dots \circ f_N$ is part of a branch of f^N), then

$$\text{var}(Q_1 P_{f_1} Q_2 P_{f_2} \dots Q_N P_{f_N} h) \leq \left((1 + \frac{3}{2}\beta)^{2N} \cdot \lambda^{-1} + \frac{1}{2} N \beta \eta^{-N} \right) \cdot \text{var}(h) + \lambda^{-1} \tilde{C}_N \cdot \|h\|_1 .$$

for each $h \in \text{BV}$.

Proof. Because of the variation condition (3.13)

$$\text{var}\left(\frac{1}{\tilde{f}'_j}\right) \leq \beta \cdot \left\| \frac{1}{\tilde{f}'_j} \right\|_\infty \leq \frac{\beta}{\eta} .$$

Therefore a simple induction yields

$$\text{var}\left(\frac{1}{|(f_1 \circ \dots \circ f_N)'|}\right) \leq N \beta \eta^{-N} .$$

Hence

$$\begin{aligned}
& \text{var}(Q_1 P_{f_1} Q_2 P_{f_2} \dots Q_N P_{f_N} h) \\
&= \text{var}(P_{\tilde{f}_1} P_{\tilde{f}_2} \dots P_{\tilde{f}_N} \tilde{Q}_N h) \\
&= \text{var}\left(\frac{\tilde{Q}_N h}{|(\tilde{f}_1 \circ \dots \circ \tilde{f}_N)'|} \circ (\tilde{f}_1 \circ \dots \circ \tilde{f}_N)^{-1}\right) \\
&\leq \text{var}(\tilde{Q}_N h) \cdot \lambda^{-1} + \underbrace{\|\tilde{Q}_N h\|_\infty}_{\leq \frac{1}{2} \text{var}(\tilde{Q}_N h)} \cdot N \beta \eta^{-N} \\
&\leq \left(\left(1 + \frac{3}{2}\beta\right)^{2N} \cdot \lambda^{-1} + \frac{1}{2} N \beta \eta^{-N} \right) \cdot \text{var}(h) + \lambda^{-1} \tilde{C}_N \cdot \|h\|_1 .
\end{aligned}$$

■

Lemma 3.7 *Let*

$$Z = Z_N \cap f_N^{-1} Z_{N-1} \cap (f_{N-1} \circ f_N)^{-1} Z_{N-2} \dots \cap (f_2 \circ \dots \circ f_N)^{-1} Z_1 ,$$

and as before $f_i = f|_{Z_i \setminus Y}$. (Note that $Z \in \mathcal{Z}_N$ if $Z \neq \emptyset$.) Recall that $\Lambda = \sup |f'| < \infty$. If

$$\text{spread}(Q) < \frac{\Lambda - 1}{\Lambda^N} \cdot \min\{|J_i| : i = 1, \dots, r\} ,$$

then

$$Q P_{f_1} \dots Q P_{f_N}(h) = Q P_{f_1} \dots Q P_{f_N}(h \cdot 1_Z)$$

for each $h \in \text{BV}$.

Proof. Let $Z' \in \mathcal{Z}_N$, $Z' \neq Z$. We prove that

$$Q P_{f_1} \dots Q P_{f_N}(h \cdot 1_{Z'}) = 0 . \quad (3.14)$$

Suppose this is not true. Then there is at least one δ -pseudoorbit $x_N, x_{N-1}, \dots, x_1, x_0$ with

- $\delta = \text{spread}(Q)$,
- $x_N \in Z'$, in particular $f^{N-j}(x_N) \in Z'_j$ ($j = N, \dots, 1$),
- $x_j \in Z_j \setminus Y$ ($j = N, \dots, 1$), in particular $Z'_N = Z_N$, and
- $|x_{j-1} - f_j(x_j)| < \delta$ ($j = N, \dots, 1$).

It follows that

$$|x_j - f^{N-j}(x_N)| < \sum_{i=0}^{N-j-1} \delta \Lambda^i < \delta \frac{\Lambda^N}{\Lambda - 1} < \min\{|J_i| : i = 1, \dots, r\}$$

for $j = N, \dots, 1$ such that $x_j \in Z'_j \cup Y$. But this is possible only if $Z_j = Z'_j$ ($j = N, \dots, 1$), i.e. if $Z = Z'$, a contradiction. ■

Proposition 3.8 *If the assumptions (3.1), (3.2), (3.3), (3.6), and (3.7) are satisfied, then there are $C_1, C_2 > 0$, $\alpha \in (0, 1)$ and $\delta > 0$ such that for all $h \in \text{BV}$ and all $k \in \mathbb{N}$*

$$\text{var}(\tilde{P}_1^k h) \leq C_1 \alpha^k \cdot \text{var}(h) + C_2 \cdot \|h\|_1$$

if $\text{spread}(Q) < \delta$ and $|J_j| < \delta$ for all $j = 1, \dots, r$. In fact, $C_1 = \left(\frac{2}{\eta}\right)^N$, $\alpha = \left(\frac{3}{4}\right)^{1/N}$, $C_2 = C_2(N, C, f)$.

Proof. Let $M : \text{BV} \rightarrow \text{BV}$ denote multiplication by $1_{X \setminus Y}$. In view of the previous lemma we can expand $\tilde{P}_1^N(h)$ as

$$\begin{aligned} \tilde{P}_1^N(h) &= (QP_f M)^N(h) = \sum_{Z_1, \dots, Z_N \in \mathcal{Z}} QP_{f_{Z_1}} M \dots QP_{f_{Z_N}} M(h) \\ &= \sum_{Z \in \mathcal{Z}_N} Q(P_{f_1} Q) \dots (P_{f_{N-1}} Q) P_{f_N}(h \cdot 1_Z). \end{aligned}$$

Hence, by Lemma 3.6

$$\begin{aligned} \text{var}(\tilde{P}_1^N(h)) &\leq \sum_{Z \in \mathcal{Z}_N} \left[\left(\left(1 + \frac{3}{2}\beta\right)^{2N} \cdot \lambda^{-1} + \frac{1}{2}N\beta\eta^{-N} \right) \cdot \text{var}(h \cdot 1_Z) + \lambda^{-1}\tilde{C}_N \cdot \|h \cdot 1_Z\|_1 \right] \\ &\leq \sum_{Z \in \mathcal{Z}_N} \left[\left(2\left(1 + \frac{3}{2}\beta\right)^{2N} \cdot \lambda^{-1} + N\beta\eta^{-N} \right) \cdot \text{var}_Z(h) \right. \\ &\quad \left. + \max\{|Z|^{-1} : Z \in \mathcal{Z}_N\} \lambda^{-1}\tilde{C}_N \cdot \|h \cdot 1_Z\|_1 \right] \\ &\leq \left(2\left(1 + \frac{3}{2}\beta\right)^{2N} \cdot \lambda^{-1} + N\beta\eta^{-N} \right) \cdot \text{var}(h) + \max\{|Z|^{-1} : Z \in \mathcal{Z}_N\} \lambda^{-1}\tilde{C}_N \cdot \|h\|_1. \end{aligned}$$

Now choose β so small that $2(1 + \frac{3}{2}\beta)^{2N} \cdot \lambda^{-1} + N\beta\eta^{-N} < (3\lambda^{-1})^{\frac{1}{N}}$ and observe that $\lambda > 4$. This proves the proposition for $k = N$ with $C_1 = 1$, $\alpha = (\frac{3}{4})^{1/N}$ and some $C_2 = C_2(N, C, f)$. Iterated application of this inequality extends it to integer multiples k of N with a new $C_2 = \frac{1}{1-\alpha}C_{2,\text{old}}$. The extension to general k follows from the elementary inequality

$$\text{var}(\tilde{P}_1 h) \leq \frac{2}{\eta} \text{var}(h \cdot 1_{X \setminus Y}) + (C + \text{Const}) \cdot \|h\|_1 \leq \frac{4}{\eta} \text{var}(h) + (C + \text{Const}) \cdot \|h\|_1$$

with a constant Const depending only on f . ■

3.2.2 Proof of Theorem 1.2

If f has no periodic turning points, then $\tilde{P}_2^k = 0$ for some $k \in \mathbb{N}$ that depends only on f , such that (3.9) is trivially satisfied for $\tilde{P}_2$ with constants C_1, C_2, α depending only on f and Q . In view of Lemma 3.3 this implies Theorem 1.2.

3.3 Estimates close to turning points

In this section we prove (3.9) for $\tilde{P}_2$, i.e. for branches with orbits close to turning points.

3.3.1 The random walk assumption

In order to prevent perturbed trajectories from being trapped in small neighbourhoods of periodic turning points (this was the case in the example from Lemma 2.2), we introduced the *random walk* assumption RW in (1.5). Here we relate it to a more technical *exponential decay* assumption.

Consider P_f and Q_ε ($\varepsilon > 0$) together with a decomposition $Q_\varepsilon P_f = \tilde{P}_1 + \tilde{P}_2 = \tilde{P}_{1,\varepsilon} + \tilde{P}_{2,\varepsilon}$ as in Section 3.1.4. Let $* \in \{L^1, L^\infty, \text{BV}\}$. We say that

$$\tilde{P}_2 \text{ satisfies } ED_*, \text{ if there are constants } C > 0 \text{ and } \omega \in (0, 1) \text{ such that } \|\tilde{P}_{2,\varepsilon}^n\|_* \leq C\omega^n \quad (3.15)$$

In order to derive ED_{L^1} from RW , the intervals J_k , whose union makes up Y , must be chosen more carefully than in (3.5) and (3.7) where we only took care that $f(J_j) \supset \tilde{J}_i$ for some i if $f(J_j) \cap \tilde{Y} \neq \emptyset$

and that random jumps from $X \setminus \tilde{Y}$ to Y are excluded. In particular, $\text{spread}(Q) \leq |fJ_j| - |J_j|$. Now we require that this inequality is in fact nearly an equality. More exactly: Let Y be the smallest domain such that

$$f_\varepsilon(B_\varepsilon(Y) \setminus Y) \cap Y = \emptyset.$$

Proposition 3.9 *If Q_ε satisfies RW and if $f \in H_\infty$, then $\tilde{P}_2$ satisfies ED_{L^1} .*

Consider a periodic turning point c with a period N . We assume first that the map f is locally linear. Let us denote

$$\lambda = \min \left\{ \prod_{k=1}^{2N} |f'(f^k x)| : x, f^N x \in B_\varepsilon(c) \right\}.$$

Lemma 3.10 *Let $f \in H_\infty$, $f^N(c) = c$, $Q_\varepsilon \in RW$. Then $\mathcal{P}\{f_\varepsilon^{2nN}(x) \notin Y\} \geq \delta^{2nN}$ for each $x \in Y$ and $n = -\ln(\theta(\lambda - 1))/\ln \theta$.*

Proof. The domain Y consists of intervals around the points of the trajectory of the periodic turning point c . Let us denote the interval around the point c by J . Then the distance from its boundary points to the point c is not more, than $\varepsilon/(\lambda - 1)$.

Note, that $\lambda > 1$. Indeed, after N iterations any point from the neighborhood of the point c either will return to the neighborhood, or will leave the interval J (if the map is discontinuous at the point c). Therefore if $\lambda \leq 1$, then the H_∞ -assumption does not hold.

Let us fix a point $x \in J$. Then, applying H_∞ and RW assumptions, we have

$$\mathcal{P}\{\text{dist}(f_\varepsilon^{2nN} x, c) > \theta \varepsilon \lambda^n\} \geq \delta^{2nN}$$

for any integer n and for any point $x \in J$. Therefore, if

$$\theta \varepsilon \lambda^n > \frac{\varepsilon}{\lambda - 1},$$

then any point will leave the interval J in $2nN$ iterations with some positive probability, which will not depend on ε . The value of n in the statement of lemma satisfies this inequality. ■

Corollary 3.11 *For small ε the conclusion of the lemma remains true for piecewise C^2 -maps, if the constant λ is replaced by a slightly smaller one.*

Now to finish the proof of the proposition, we need the following simple statement.

Lemma 3.12 *Let Q be any transition kernel satisfying $Q(x, X \setminus Y) \geq \delta$ for any point $x \in Y$. Then for any nonnegative function $h \in L^1$ with the support in the set Y we have:*

$$\int_Y Qh(x) dx \leq (1 - \delta) \int_Y h(x) dx.$$

Proof. Recall that for any measurable set A we have

$$\int_A Qh(x) dx = \int Q(x, A)h(x) dx.$$

Thus

$$\int_Y Qh(x) dx = \int Q(x, Y)h(x) dx \leq (1 - \delta) \int_Y h(x) dx.$$

■

3.3.2 Smooth perturbations

We start with a special case of absolutely continuous perturbations for which the calculations are particularly easy.

Proposition 3.13 *Suppose that Q is an integral operator with kernel $q(x, y)$ satisfying*

$$\text{var}(y \mapsto q(x, y)) \leq \frac{M}{|Y|} \quad \text{for all } x$$

and such that $\tilde{P}_2$ satisfies ED_{L^1} . Then

$$\text{var}(\tilde{P}_2^n h) \leq MC\omega^{n-1} \cdot \left(\text{var}(h) + \frac{1}{|X|} \|h\|_1 \right)$$

for $h \in \text{BV}$.

Proof. Let $h \in \text{BV}$ and denote $\tilde{h} := P(h1_Y)$. Then $\tilde{P}_2 h = Q\tilde{h}$ and

$$\begin{aligned} \text{var}(\tilde{P}_2 h) &= \text{var} \left(y \mapsto \int \tilde{h}(x) q(x, y) dx \right) \\ &= \sup_{\varphi \in \mathcal{T}_1(X)} \int \int \tilde{h}(x) \varphi'(y) q(x, y) dx dy \\ &\leq \int |\tilde{h}(x)| \cdot \left| \sup_{\varphi \in \mathcal{T}_1(X)} \int \varphi'(y) q(x, y) dy \right| dx \\ &\leq \int |\tilde{h}(x)| \cdot \text{var}(y \mapsto q(x, y)) dx \\ &\leq \frac{M}{|Y|} \int |P(h1_Y)(x)| dx \\ &\leq \frac{M}{|Y|} \int_Y |h(x)| dx. \end{aligned}$$

As $\tilde{P}_2 h = QP_f(h1_Y) = \tilde{P}_2(h1_Y)$, it follows that

$$\begin{aligned} \text{var}(\tilde{P}_2^n h) &\leq \frac{M}{|Y|} \int |\tilde{P}_2^{n-1} h(x)| dx \\ &\leq \frac{M}{|Y|} \int \tilde{P}_2^{n-1} |h(x)1_Y(x)| dx \\ &\leq \frac{M}{|Y|} \cdot \|\tilde{P}_2^{n-1}\|_1 \cdot \int_Y |h(x)| dx \\ &\leq MC\omega^{n-1} \cdot \|h\|_\infty \\ &\leq MC\omega^{n-1} \cdot \left(\text{var}(h) + \frac{1}{|X|} \|h\|_1 \right) \end{aligned}$$

■

Theorem 1.4 follows from this proposition and Proposition 3.9 via Lemma 3.3.

3.3.3 General perturbations

We now turn to more general perturbations, which are much more delicate. Our first goal is to decompose the operator $\tilde{P}_2$ as $\tilde{P}_2 = L + \tilde{P}_2 K$, where

$$\text{var}_Y(L^N h) \leq \alpha \cdot \text{var}_Y(h) \quad \text{for } h \in \text{BV}(Y)$$

with some suitable constant $0 < \alpha < 1$ and where K is a finite rank operator such that also $\tilde{P}_2 K$ is of finite rank. Our choice of K is

$$Kh = \sum_{j=1}^r h(c_j) 1_{J_j} .$$

As our BV-functions are in fact L^1 -equivalence classes, one should interpret $h(c_j)$ as $\frac{1}{2}(\text{ess-lim}_{x \nearrow c_j} h(x) + \text{ess-lim}_{x \searrow c_j} h(x))$. For later use we note that

$$L(h \cdot 1_{X \setminus Y}) = \tilde{P}_2(h \cdot 1_{X \setminus Y}) - \sum_{j=1}^r h(c_j) \cdot \tilde{P}_2 1_{J_j} = 0 .$$

For the formulation of the next lemma we introduce the following notation: Let $A, B \subseteq \{1, \dots, r\}$. Then

$$A \rightarrow B \quad \text{if} \quad \exists i \in A, j \in B : f(c_i) = c_j .$$

For $i_0 \in \{1, \dots, r\}$ let

$$\begin{aligned} \Lambda_N(i_0) &:= \min \left\{ \prod_{l=0}^{N-1} |f'(c_{i_l})| : i_0 \rightarrow \{i_1, \hat{i}_1\}, \{i_l, \hat{i}_l\} \rightarrow \{i_{l+1}, \hat{i}_{l+1}\} \ (l = 1, \dots, N-1) \right\} \\ \Lambda_N &:= \min \{ \Lambda_N(i_0) : i_0 \in \{1, \dots, r\} \} \end{aligned}$$

and observe that

$$\Lambda_{N+1}(i_0) = |f'(c_{i_0})| \cdot \min \left\{ \Lambda_N(i_1) : i_0 \rightarrow \{i_1, \hat{i}_1\} \right\} \quad (3.16)$$

Lemma 3.14 *Fix $N \in \mathbb{N}$ and $\kappa > 1$. There is $\delta > 0$ depending only on f and C such that for a refined partition $\mathcal{Z}$ with $\text{diam}(\mathcal{Z}) < \delta$ holds:*

$$\text{var}_Y(L^N h) \leq \sum_{j=1}^r \text{var}_Y(L^N(h \cdot 1_{J_j})) \leq \kappa \Lambda_N^{-1} \cdot \text{var}_Y(h) .$$

Proof. For $j, k \in \{1, \dots, r\}$

$$\begin{aligned} \text{var}_{J_k \cup J_{\hat{k}}}(L(h 1_{J_j})) &= \text{var}_{J_k \cup J_{\hat{k}}}(Q P_f((h - h(c_j)) \cdot 1_{J_j})) \\ &\leq \text{var}_{\tilde{J}_k \cup \tilde{J}_{\hat{k}}}(P_f[(h - h(c_j)) \cdot 1_{J_j}]) + C \cdot \|P_f[(h - h(c_j)) \cdot 1_{J_j}] \cdot 1_{\tilde{J}_k \cup \tilde{J}_{\hat{k}}}\|_1 \\ &= \text{var}_{\tilde{J}_k \cup \tilde{J}_{\hat{k}}} \left(\frac{h - h(c_j)}{|f'|} \circ f_j^{-1} \cdot 1_{f J_j} \right) + C \cdot \|(h - h(c_j)) \cdot 1_{J_j} \cdot 1_{\tilde{J}_k \cup \tilde{J}_{\hat{k}}}\|_1 \end{aligned}$$

by (3.6) and the fact that P_f is a L^1 -contraction. By (3.5) we have either $f(J_j) \cap (\tilde{J}_k \cup \tilde{J}_{\hat{k}}) = \emptyset$ such that $j \not\rightarrow \{k, \hat{k}\}$ or $f(J_j) \supset \tilde{J}_k$ or $\supset \tilde{J}_{\hat{k}}$ in such a way that $j \rightarrow \{k, \hat{k}\}$ but $f(d_j) \notin \tilde{J}_k \cup \tilde{J}_{\hat{k}}$. In the first case

$$\text{var}_{J_k \cup J_{\hat{k}}}(L(h \cdot 1_J)) = 0 .$$

In the second case $f(d_j) \notin \tilde{J}_k \cup \tilde{J}_{\hat{k}}$ and $\frac{h(x) - h(c_j)}{|f'|} \Big|_{x=c_j} = 0$, whence

$$\begin{aligned} &\text{var}_{\tilde{J}_k \cup \tilde{J}_{\hat{k}}} \left(\frac{h - h(c_j)}{|f'|} \circ f_j^{-1} \cdot 1_{f J_j} \right) = \text{var}_{J_j} \left(\frac{h - h(c_j)}{|f'|} \right) \\ &= \text{var}_{J_j} \left(\frac{h - h(c_j)}{|f'(c_j)|} \frac{|f'(c_j)|}{|f'|} \right) \leq \text{var}_{J_j}(h - h(c_j)) \cdot \frac{1 + \beta}{|f'(c_j)|} + \sup_{J_j} (h - h(c_j)) \cdot \beta \\ &\leq \text{var}_{J_j}(h) \cdot \left(\frac{1 + \beta}{|f'(c_j)|} + \beta \right) \end{aligned}$$

by the assumptions on β in (3.3). Hence

$$\text{var}_{J_k \cup J_{\hat{k}}}(L(h1_{J_j})) \leq \text{var}_{J_j}(h) \cdot \left(\frac{1+\beta}{|f'(c_j)|} + \beta + C \cdot |J_j| \text{var}_{J_j}(h) \right) \cdot \delta_{j \rightarrow \{k, \hat{k}\}}$$

Let $\eta_0 = \min\{\eta, 1\}$. Given $N > 0$ and choosing $\mathcal{Z}$ sufficiently fine we can make $\beta > 0$ and $|J_j|$ so small that

$$\frac{1+\beta}{|f'(c_j)|} + \beta < \frac{1+\frac{\varepsilon}{2}}{|f'(c_j)|}, \quad C \cdot |J_j| < \frac{\varepsilon \eta_0^{2N}}{|f'(c_j)|} \quad \text{and} \quad 1+\varepsilon < \kappa^{1/N}$$

for a suitable $\varepsilon > 0$ and all $j = 1, \dots, r$. (Observe that r , the cardinality of $\mathcal{Z}$, may depend on the refinement of $\mathcal{Z}$.) Hence

$$\text{var}_{J_k \cup J_{\hat{k}}}(L(h1_{J_j})) \leq \left(\frac{1+\frac{\varepsilon}{2}}{|f'(c_j)|} + \frac{\varepsilon \eta_0^{2N}}{|f'(c_j)|} \right) \cdot \delta_{j \rightarrow \{k, \hat{k}\}} \cdot \text{var}_{J_j}(h) \quad (3.17)$$

for all $j = 1, \dots, r$, and summation over k yields

$$\begin{aligned} \text{var}_Y(L(h1_{J_j})) &= \frac{1}{2} \sum_{k=1}^r \text{var}_{J_k \cup J_{\hat{k}}}(L(h1_{J_j})) \leq \left(\frac{1+\frac{\varepsilon}{2}}{|f'(c_j)|} + \frac{\varepsilon}{|f'(c_j)|} \right) \cdot \text{var}_{J_j}(h) \\ &\leq \frac{\kappa^{1/N}}{|f'(c_j)|} \cdot \text{var}_{J_j}(h) \end{aligned}$$

for all $j = 1, \dots, r$. This is the special case $n = 1$ of the estimate

$$\text{var}_Y(L^n(h1_{J_j})) \leq \frac{\kappa^{n/N}}{\Lambda_n(j)} \cdot \text{var}_{J_j}(h) \quad (3.18)$$

which we now prove for general n inductively: Suppose (3.18) holds for $n \in \{1, \dots, N-1\}$. Then

$$\begin{aligned} \text{var}_Y(L^{n+1}(h1_{J_j})) &= \sum_{k=1}^r \text{var}_Y(L^n(L(h1_{J_j})1_{J_k})) \\ &\leq \sum_{k=1}^r \frac{\kappa^{n/N}}{\Lambda_n(k)} \cdot \text{var}_{J_k}(L(h1_{J_j})) \\ &= \frac{1}{2} \sum_{k=1}^r \frac{\kappa^{n/N}}{\Lambda_n(k)} \cdot \text{var}_{J_k \cup J_{\hat{k}}}(L(h1_{J_j})) \\ &\leq \frac{1}{2} \sum_{k=1}^r \frac{\kappa^{n/N}}{\Lambda_n(k)} \cdot \left(\frac{1+\frac{\varepsilon}{2}}{|f'(c_j)|} \cdot \delta_{j \rightarrow \{k, \hat{k}\}} + \frac{\varepsilon \eta_0^{2N}}{r|f'(c_j)|} \right) \cdot \text{var}_{J_j}(h) \\ &\leq \kappa^{n/N} \left(\frac{1+\frac{\varepsilon}{2}}{\Lambda_{n+1}(j)} + \frac{\varepsilon \eta_0^N}{2} \right) \cdot \text{var}_{J_j}(h) \\ &\leq \kappa^{n/N} \frac{1+\varepsilon}{\Lambda_{n+1}(j)} \cdot \text{var}_{J_j}(h) \\ &\leq \frac{\kappa^{(n+1)/N}}{\Lambda_{n+1}(j)} \cdot \text{var}_{J_j}(h) \end{aligned}$$

by (3.17) and (3.16). This proves (3.18) for $n = N$ in particular. Summing over all j finally yields the claim of the lemma. ■

Corollary 3.15 *If there is some N such that $\Lambda_N > 1$, then there are $C(\Lambda_1, N) > 0$ depending continuously on Λ_1 and $\delta > 0$ depending only on f and C such that for a partition $\mathcal{Z}$ with $\text{diam}(\mathcal{Z}) < \delta$ and $\alpha := \Lambda_N^{-\frac{1}{2N}}$ holds*

$$\text{var}_Y(L^n h) \leq \sum_{j=1}^r \text{var}_Y(L^n(h \cdot 1_{J_j})) \leq C(\Lambda_1, N) \cdot \alpha^n \cdot \text{var}_Y(h)$$

for all $n > 0$.

Proof. Let $\kappa = \Lambda_N^{\frac{1}{2}}$. Then the estimate with $C = 1$ follows for integer multiples n of N from the previous lemma. To pass from this to general n , apply the previous lemma successively with $N = 1$. ■

We turn to the finite-dimensional contribution $\tilde{P}_2 K$ of $\tilde{P}_2$: Two straightforward inductions yield

$$\tilde{P}_2^n = \sum_{k=0}^{n-1} \tilde{P}_2^{n-k} K L^k + L^n \quad \text{and} \quad (3.19)$$

$$\tilde{P}_2^n = \sum_{k=0}^{n-1} L^k \tilde{P}_2 K \tilde{P}_2^{n-k-1} + L^n. \quad (3.20)$$

Inserting (3.20) into (3.19) results in

$$\tilde{P}_2^n = \sum_{k=0}^{n-1} \sum_{l=0}^{n-k-1} L^l \tilde{P}_2 K \tilde{P}_2^{n-k-l-1} K L^k + \sum_{k=0}^{n-1} L^{n-k} K L^k + L^n$$

whence

$$\begin{aligned} \tilde{P}_2^n h &= \sum_{k=0}^{n-1} \sum_{l=0}^{n-k-1} \sum_{i=1}^r \sum_{j=1}^r L^l (\tilde{P}_2 1_{J_j}) \cdot (\tilde{P}_2^{n-k-l-1} 1_{J_i})(c_j) \cdot (L^k h)(c_i) \\ &\quad + \sum_{k=0}^{n-1} \sum_{i=1}^r (L^{n-k} 1_{J_i}) \cdot (L^k h)(c_i) + L^n h \end{aligned} \quad (3.21)$$

Observe now that for $k \geq 1$

$$L^k h = \sum_{i=1}^r L(L^{k-1} h \cdot 1_{J_i}) = \sum_{i=1}^r \tilde{P}_2 v_i \quad (3.22)$$

where $v_i = (K - \text{Id})(L^{k-1} h \cdot 1_{J_i})$ such that $\text{supp}(v_i) = J_i$ and $v_i(c_i) = 0$. Therefore

$$\begin{aligned} \text{var}(\tilde{P}_2 v_i) &= \text{var}(QP(v_i \cdot 1_Y)) \\ &\leq \text{var}(P(v_i \cdot 1_Y)) + C \cdot \|P(v_i \cdot 1_Y)\|_1 \\ &\leq \text{var}\left(\frac{v_i}{|f'|} \circ f|_{J_i}^{-1} \cdot 1_{f J_i}\right) + C \cdot \|v_i\|_1 \\ &= \text{var}\left(\frac{v_i}{|f'|} \cdot 1_{J_i}\right) + C \cdot \|v_i\|_1 \\ &= 2 \text{var}_{J_i}\left(\frac{v_i}{|f'|}\right) + C \cdot \|v_i\|_1 \quad \text{as } v_i(c_i) = 0, \\ &\leq \frac{2}{\eta} \text{var}_{J_i}(v_i) + (C + \text{Const}) \cdot \|v_i\|_1 \\ &\leq \frac{2}{\eta} \text{var}_{J_i}(L^{k-1} h) + (C + \text{Const}) \cdot \|v_i\|_\infty \cdot |J_i| \\ &\leq \left(\frac{2}{\eta} + (C + \text{Const}) \cdot |J_i|\right) \cdot \text{var}_{J_i}(L^{k-1} h) \end{aligned}$$

with a constant Const depending only on f . It follows that for sufficiently small J_i (depending on f and C)

$$\begin{aligned} 2\|L^k h\|_\infty &\leq \text{var}(L^k h) \\ &\leq \sum_{i=1}^r \text{var}(\tilde{P}_2 v_i) \leq \frac{3}{\eta} \cdot \text{var}_Y(L^{k-1} h) \\ &\leq \frac{3}{\eta} \cdot C(\Lambda_1, N) \cdot \alpha^{k-1} \cdot \text{var}(h) . \end{aligned} \quad (3.23)$$

As $|h(c_i)| \leq \frac{1}{2} \text{var}(h)$, we can assume that this inequality holds for $k = 0$, too. Now (3.21) implies

$$\begin{aligned} &\text{var}(\tilde{P}_2^n h) \\ &\leq \frac{3}{\eta} C(\Lambda_1, N)^2 n \sum_{i=1}^r \left[\sum_{j=1}^r \left(\text{var}(\tilde{P}_2 1_{J_j}) \cdot \sum_{s=0}^{n-1} \alpha^{s-1} |(\tilde{P}_2^{n-s-1} 1_{J_i})(c_j)| \right) + \alpha^{n-1} \right] \cdot \text{var}(h) , \end{aligned}$$

and the problem of proving exponential convergence of $\text{var}(\tilde{P}_2^n h)$ to 0 is reduced to proving exponential convergence of $(\tilde{P}_2^n 1_{J_i})(c_j)$ to 0. As $\text{var}(\tilde{P}_2 1_{J_j}) \leq \frac{2}{\eta} + (C + \text{Const}) \cdot |J_j|$ with a constant depending only on f , this proves

Proposition 3.16 *Suppose there are constants $B, \beta > 0$ such that for all $i, j = 1, \dots, r$*

$$|(\tilde{P}_2^n 1_{J_i})(c_j)| \leq B \cdot \beta^n . \quad (3.24)$$

Let α be the constant from Corollary 3.15, $\hat{\alpha} := \max\{\alpha, \beta\}$. Then there are constants $C_2, \delta_2 > 0$ depending only on f, C and B and such that for a refined partition $\mathcal{Z}$ with $\text{diam}(\mathcal{Z}) < \delta_2$ holds

$$\text{var}(\tilde{P}_2^n h) \leq C_2 \cdot \hat{\alpha}^n \cdot \text{var}(h) \quad \text{for all } h \in \text{BV}.$$

Unfortunately assumption (3.24) does not follow automatically from RW or ED_{L^1} . Instead it is implied by the stronger assumption ED_{L^∞} . Therefore we discuss the following approach. By inequality (3.23),

$$\text{var}(L^k h) \leq \text{Const} \cdot \alpha^k \cdot \text{var}(h) .$$

As $\text{supp}(Lh) \subseteq \tilde{Y}$ for each $h \in \text{BV}$, L leaves $\text{BV}(\tilde{Y})$ invariant, and $\text{var} = \text{var}_X$ is a norm on $\text{BV}(\tilde{Y})$. $\tilde{K}$ is a bounded finite-rank operator on $\text{BV}(\tilde{Y})$, because the evaluation of a function h at one point is a bounded linear functional. In particular, also $\tilde{K}_n := \tilde{P}_2^n - L^n = (L + \tilde{K})^n - L^n$ has finite rank, and as

$$\text{var}((\tilde{P}_2^n - \tilde{K}_n)h) = \text{var}(L^n h) \leq \text{Const} \cdot \alpha^n \cdot \text{var}(h) ,$$

$\tilde{P}_2$ is quasicompact as operator on $\text{BV}(\tilde{Y})$. As $\tilde{P}_2$ is at the same time a positive L^1 -operator, it has a nonnegative eigenfunction h_0 to its leading positive eigenvalue r_0 . Hence

$$r_0^n \int h_0 dm = \int \tilde{P}_2^n h_0 dm \leq \text{Const} \cdot \omega^n \int h_0 dm \quad (3.25)$$

for some $\omega \in (0, 1)$ by the ED_{L^1} assumption, which implies $r_0 \leq \omega < 1$.

Therefore $\text{var}_Y(\tilde{P}_2^{n-1} h) \leq \text{Const} \cdot \omega^n \text{var}(h)$, which is the desired estimate except that the constant, which comes from the spectral representation of $\tilde{P}_2$ cannot be controlled uniformly in ε . Nevertheless it shows that for each fixed ε the perturbed system f_ε has a smooth invariant measure, and its transition operator satisfies the Lasota-Yorke type inequality.

If f is linear in neighbourhoods of the turning points c_i and if $Q_\varepsilon(x, A) = Q(\varepsilon^{-1}(A - x))$ for a fixed probability measure Q (i.e. the usual scaling behaviour), then the constant in (3.25) is uniform in Q_ε , because the variation of a function is not changed by a linear change of scale. This finishes the **proof of Theorem 1.5**. ■

3.4 Proof of Ulam's conjecture

Another problem related to stochastic stability is the approximation of chaotic dynamics by finite state Markov chains. The general idea here is due to Ulam [18]. Consider a partition of the phase space X into a finite number of disjoint components $\{\Delta_i\}_{i=1}^K$, $0 < \theta\varepsilon \leq |\Delta_i| \leq \varepsilon$. Then one can compare statistical properties of a map f with those of a Markov chain with K states, whose transition probabilities are defined as follows:

$$p_{ij} = \frac{|f^{-1}\Delta_j \cap \Delta_i|}{|\Delta_i|}$$

Ulam's conjecture states that if one consider a sequence of finite approximations of this type with $\max |\Delta_i| \rightarrow 0$ and $K \rightarrow \infty$, invariant measures of these Markov chains converge weakly to a SBR measures of the map f . The connection of this problem to the question of stochastic stability is straightforward, because Ulam's approximation corresponds to a specific Markov random perturbation where the transition probability to go from a point $x \in X$ to a measurable set $A \subset X$ is equal to

$$Q_\varepsilon(x, A) := \sum_i \mathbf{1}_{\Delta_i}(x) \frac{|A \cap \Delta_i|}{|\Delta_i|}, \quad (3.26)$$

while the corresponding transition operator $Q_\varepsilon : L^1 \rightarrow L^1$ is defined by

$$Q_\varepsilon h(x) := \sum_i \mathbf{1}_{\Delta_i}(x) \frac{1}{|\Delta_i|} \int_{\Delta_i} h(s) ds. \quad (3.27)$$

Therefore Ulam's conjecture can be treated in the context of random perturbations and it was proved earlier for the case $\lambda_f > 2$ in [14] (see also discussion of this question in [10, 11, 2, 3]).

Let L_ε^1 be the finite dimensional linear subspace of L^1 generated by $\{\frac{1}{|\Delta_i|} \mathbf{1}_{\Delta_i}\}$. Observe that $Q_\varepsilon L^1 = L_\varepsilon^1$. Consider the transition operator $P_\varepsilon : L_\varepsilon^1 \rightarrow L_\varepsilon^1$, defined by

$$P_\varepsilon \left(\frac{1}{|\Delta_i|} \mathbf{1}_{\Delta_i} \right) := \sum_j p_{ij} \frac{1}{|\Delta_j|} \mathbf{1}_{\Delta_j}.$$

Lemma 3.17 $P_\varepsilon h = Q_\varepsilon P_f h$ for any function $h \in L_\varepsilon^1$.

Proof. It is enough to prove this statement for $h = \mathbf{1}_{\Delta_i}$. As $\int_{\Delta_j} P_f \mathbf{1}_{\Delta_i}(s) ds = |f^{-1}\Delta_j \cap \Delta_i|$, we have

$$\begin{aligned} Q_\varepsilon P_f \mathbf{1}_{\Delta_i}(x) &= \sum_j \frac{1}{|\Delta_j|} \int_{\Delta_j} P_f \mathbf{1}_{\Delta_i} \mathbf{1}_{\Delta_j}(x) = \sum_j \frac{|\Delta_i \cap f^{-1}\Delta_j|}{|\Delta_j|} \mathbf{1}_{\Delta_j}(x) \\ &= \sum_j \frac{|\Delta_i \cap f^{-1}\Delta_j|}{|\Delta_i|} \frac{|\Delta_i|}{|\Delta_j|} \mathbf{1}_{\Delta_j}(x) = |\Delta_i| \sum_j p_{ij} \frac{1}{|\Delta_j|} \mathbf{1}_{\Delta_j}(x). \end{aligned}$$

Thus

$$Q_\varepsilon P_f \left(\frac{1}{|\Delta_i|} \mathbf{1}_{\Delta_i}(x) \right) = P_\varepsilon \left(\frac{1}{|\Delta_i|} \mathbf{1}_{\Delta_i}(x) \right).$$

■

Notice that a related statement was proved in [14] for the case when all intervals Δ_i have the same length.

Straightforward calculations show that the transition probability, defined by (3.26), is smooth and satisfies the variation assumption. Actually it satisfies the stronger assumption: $\text{var}(Q_\varepsilon h) \leq \text{var}(h)$ for any $h \in \text{BV}$.

If $\text{PTP} = \emptyset$, then Ulam's conjecture is a corollary to Theorem 1.2. However, if $\text{PTP} \neq \emptyset$, formally the statement of our Theorem 1.4 can not be applied here, because the RW -assumption may not be satisfied in Ulam's case. Here we prove that under the H_∞ -assumption Ulam's construction satisfies the ED_{L^1} -condition, such that **Theorem 1.6** follows (just as Theorem 1.4) from Proposition 3.9 via Lemma 3.3.

Consider a periodic turning point c with period N . We consider only small values of ε and therefore, as in the proof of Proposition 3.9, we may assume here that the map f is locally linear. Denote

$$\lambda := \min\left\{\prod_{k=1}^{2N} |f'(f^k x)| : x, f^N x \in B_\varepsilon(c)\right\};$$

$$\lambda_k := |f'(f^k c)|, \text{ and } \Lambda_f := \max_x |f'x|.$$

Note, that due to (3.26) the escape rate (from some interval around a turning point) for any point is the same, as for the end-points of the corresponding Ulam's interval. Define the neighborhood Y of the trajectory of the periodic point c as in the Section 3.3.1.

Lemma 3.18 *Let $f \in H_\infty$, $f^N(c) = c$ and the perturbation Q_ε be defined by (3.27). Then there exist constants $\delta > 0$ and $n < \infty$ such that for sufficiently small $\varepsilon > 0$*

$$\mathcal{P}\{f_\varepsilon^{2nN}(x) \notin Y\} \geq \delta^{2nN} \quad \text{for each } x \in J$$

where J is the component of Y containing c .

Proof. In the same way as in the proof of Lemma 3.10 consider the interval J around the point c . In this case it consists of a finite number K_0 of Ulam intervals. Note that this number is uniformly bounded for all $\varepsilon > 0$.

Fix a small positive number $\sigma := \theta\varepsilon\gamma/4$, where $\gamma := \frac{\lambda(\lambda-1)}{2N\Lambda_f^{2N}}$. Let Δ_0 be the Ulam interval containing the point c . We set $A := [c, 1] \cap J$ if $|\Delta_0 \cap [c, 1]| > |\Delta_0|/2$, and $A := [0, c] \cap J$ otherwise. Define intervals A_n inductively by:

$$A_0 := \Delta_0 \cap A$$

$$A_{n+1} := f^{n+1}A \cap \tilde{A}_{n+1}, \text{ where } \tilde{A}_{n+1} = (\cup_{i: |\Delta_i \cap fA_n| > \sigma} \Delta_i).$$

Let $n_0 = \max\{n : A_n \subseteq Y\}$. Then $\tilde{A}_n \subseteq Y$ for $n = 0, \dots, n_0$, and it follows by induction that

- A_n is an interval adjacent to a point from the orbit of c for $n = 0, \dots, n_0 + 1$ and
- A_n is a union of complete Ulam intervals for $n = 0, \dots, n_0$.

Hence

$$|A_{n+1}| \geq |fA_n| - \sigma \quad \text{for } n = 0, \dots, n_0,$$

and a bit more complex calculation yields the following lower bound

$$P_\varepsilon \mathbf{1}_{A_n}(x) \geq \frac{\gamma}{8\lambda_n} \geq \frac{\gamma}{8\Lambda_f} > 0 \quad \text{for any } x \in A_{n+1}. \quad (3.28)$$

Indeed, for any Ulam interval Δ with $\Delta \cap A_{n+1} \neq \emptyset$ there exists an Ulam interval Δ' such that $\Delta' \cap A_n \neq \emptyset$ and $|\Delta \cap f\Delta'| > \min\{\frac{\sigma}{2}, \Lambda_f\theta\varepsilon\} = \frac{\sigma}{2}$. Therefore the transition probability $p_{\Delta', \Delta}$ to go from Δ' to Δ can be estimated as follows:

$$p_{\Delta', \Delta} := \frac{|\Delta' \cap f^{-1}\Delta|}{|\Delta'|} \geq \frac{1}{\lambda_n} \cdot \frac{|f\Delta' \cap \Delta|}{|\Delta'|} \geq \frac{|\Delta|}{|\Delta'|} \cdot \frac{\sigma}{2\lambda_n\theta\varepsilon} = \frac{|\Delta|}{|\Delta'|} \cdot \frac{\gamma}{8\lambda_n},$$

such that

$$P_\varepsilon \mathbf{1}_{A_n} \geq p_{\Delta', \Delta} \cdot \frac{|\Delta'|}{|\Delta|} \cdot \mathbf{1}_\Delta \geq \frac{\gamma}{8\lambda_n}.$$

Iterating the estimate (3.4) we obtain

$$|A_n| \geq \lambda_n \lambda_{n-1} \dots \lambda_1 \theta \varepsilon / 2 - \sigma (\lambda_n \lambda_{n-1} \dots \lambda_2 + \lambda_n \lambda_{n-1} \dots \lambda_3 + \dots + \lambda_n),$$

where the term

$$z_n := \lambda_n \lambda_{n-1} \dots \lambda_2 + \lambda_n \lambda_{n-1} \dots \lambda_3 + \dots + \lambda_n$$

can be estimated from above as follows. The sequence $\{\lambda_k\}$ is periodic with period $2N$ and the product $\prod_{k=1}^{2N} \lambda_k = \lambda > 1$. Therefore the highest order term contributes most, which gives the following estimate:

$$z_{2kN} \leq \lambda^{k-1} \frac{2N \Lambda_f^{2N}}{(\lambda - 1)}.$$

Thus for $2kN \leq n_0 + 1$ holds

$$|A_{2kN}| \geq \lambda^k \left(\frac{\theta \varepsilon}{2} - \frac{2N \Lambda_f^{2N}}{\lambda(\lambda - 1)} \sigma \right) \geq \lambda^k \frac{\theta \varepsilon}{4},$$

which gives an exponential expansion rate for lengths of the intervals A_n . On the other hand by (3.28)

$$P_\varepsilon^{2kN} \mathbf{1}_{A_0}(x) \geq \left(\frac{\gamma}{8\Lambda_f} \right)^{2kN}$$

for each point $x \in A_{2kN}$. This finishes the proof because the last two inequalities provide a uniform estimate on the necessary number of steps and the probability to leave Y starting from the “centre” interval Δ_0 of Ulam’s partition. ■

Now the ED_{L^1} -property for the operator $\tilde{P}_2$ corresponding to the Ulam P_ε follows as in Proposition 3.9 via Lemma 3.12.

Remark, that if Ulam’s perturbations are selfsimilar for all $\varepsilon > 0$ (this means that the corresponding transition probabilities do not depend on ε and that f is piecewise linear), then one can prove the convergence of invariant measures in a much more simple way. Indeed, if the map f is topologically mixing, then the same is true for the corresponding finite Markov chains in Ulam’s construction. Now, due to the fact that the transition probabilities are the same for any $\varepsilon > 0$, we deduce that rates of convergence are also the same, which gives the desired statement.

Dealing with Ulam’s conjecture, we assume that the elements Δ_i of the partition are of comparable size ($0 < \theta \leq |\Delta_i|/|\Delta_j| \leq 1/\theta$). To show that this assumption is necessary, consider the simplest case of a map f with fixed turning point $c = f(c)$ and a family $\{\Delta_i^\varepsilon\}_\varepsilon$ of Ulam intervals, such that the intervals near the point c (in the interval $J = J(\varepsilon)$) are of length ε^2 , while the others are of order ε . Then the number of intervals Δ_i^ε which have nonempty intersection with J is of order $1/\varepsilon$, and for any fixed $\delta > 0$ the number of steps for a point close to c to leave the interval J with the probability at least δ goes to infinity as $\varepsilon \rightarrow 0$.

Note that Ulam’s conjecture seems quite general, and actually we do not know any counterexample even for multidimensional hyperbolic maps or maps with singularities. For example, the conjecture clearly is true for nonchaotic maps with stable periodic orbits.

3.5 The variation condition on Q

In this section we give sufficient conditions for the Lasota-Yorke-type property (1.3) and (3.6) of Q . Recall the notations from Section 3.1.2 on the stochastic part of the dynamics where we discussed the operator Q and its dual Q^* , both defined in terms of the (sub)-Markovian transition kernel $Q(x, A)$. Extending $Q(x, A)$ to all $x \in \mathbb{R}$ by setting $Q(x, A) = 0$ for $x \in \mathbb{R} \setminus X$, we can assume that Q and Q^* act on $L_m^1(\mathbb{R})$ and $L_m^\infty(\mathbb{R})$ respectively.

Let $V = X$, $\tilde{V} = \mathbb{R}$ or $V = J_k \cup J_{\hat{k}}$, $\tilde{V} = \tilde{J}_k \cup \tilde{J}_{\hat{k}}$, and denote by a and b the left and right endpoints of $\tilde{V}$ respectively. Then $V \subseteq (a + \text{spread}(Q), b - \text{spread}(Q))$ such that

$$Q^* 1_{(a,x]}(a) = 1 \text{ and } Q^* 1_{(a,x]}(b) = 0 \text{ for each } x \in V. \quad (3.29)$$

Proposition 3.19 *Suppose Q can be decomposed as a sum of linear operators $Q = R + S$ in such a way that*

$$\begin{aligned} R 1_{\tilde{V}} &= 1 \text{ on } V \text{ and } R 1_W = 0 \text{ on } V \text{ if } W \cap \tilde{V} = \emptyset, \\ \alpha_R &:= \sup_t \text{var}_V(R 1_{(a,t]}) < \infty, \\ Sh(y) &= \int h(x) s(x, y) m(dx) \text{ for some kernel } s \text{ with } s(x, y) = 0 \text{ if } x \notin \tilde{V}, y \in V, \text{ and} \\ C_S &:= \sup_x \text{var}_y s(x, y) < \infty. \end{aligned}$$

Then

$$\text{var}_V(Qh) \leq \alpha_R \cdot \text{var}_{\tilde{V}}(h) + C_S \cdot \|h \cdot 1_{\tilde{V}}\|_1 \quad (3.30)$$

for each $h \in L^1$.

Proof. For $\varphi \in \mathcal{T}_1(V)$ and $t \in X$ let $\Phi(t) := \int_a^t R^* \varphi' dm$. Then $\Phi' = R^* \varphi'$ and

$$\Phi(t) = \int R 1_{(a,t]} \cdot \varphi' dm,$$

whence $\Phi(t) = 0$ for $t \leq a$, and for $t \geq b$ holds:

$$\Phi(t) = \int R 1_{(a,t]} \varphi' dm = \int_V (R 1_{\tilde{V}} + R 1_{(a,t] \setminus \tilde{V}}) \cdot \varphi' dm = \int_V \varphi' dm = \varphi(b) - \varphi(a) = 0,$$

because $\text{supp}(\varphi') \subseteq V$. Furthermore

$$|\Phi(t)| \leq \text{var}_V(R 1_{(a,t]}) \leq \alpha_R.$$

Therefore $\alpha_R^{-1} \Phi \in \mathcal{T}_1(\tilde{V})$ and

$$\begin{aligned} \int \varphi' Qh dm &= \int \varphi' Rh dm + \int \varphi' Sh dm \\ &= \int R^* \varphi' h dm + \int \int \varphi'(y) h(x) s(x, y) m(dx) m(dy) \\ &= \int \Phi' h dm + \int \left[h(x) 1_{\tilde{V}}(x) \cdot \int s(x, y) \varphi'(y) m(dy) \right] m(dx) \\ &\leq \alpha_R \cdot \text{var}_{\tilde{V}}(h) + C_S \cdot \|h \cdot 1_{\tilde{V}}\|_1. \end{aligned}$$

■

Corollary 3.20 *The assumptions of the previous proposition are satisfied in each of the following two situations:*

1. *Q is a bistochastic kernel (i.e. $Q1 = 1$) and $Q1_{(a,x]}(y)$ is decreasing as a function of y for each fixed x (i.e. the “probability” to reach y from $(a,x]$ is decreasing in y). In this case $R = Q$ and $S = 0$ such that $\alpha_R = 1$ and $C_S = 0$.*
2. *Q has a differentiable transition density $q(x, y)$ and*

$$C_q := \sup_x \int \left| \frac{\partial q}{\partial x}(x, y) + \frac{\partial q}{\partial y}(x, y) \right| m(dy) < \infty . \quad (3.31)$$

In this case $\alpha_R = 1$ and $C_S = C_q$.

Proof.

1. As $Q(x, V) = 0$ for $x \notin \tilde{V}$, we have $Q1_W = 0$ on V if $W \cap \tilde{V} = \emptyset$ and $Q1_{\tilde{V}} = Q1 - Q1_{\mathbb{R} \setminus \tilde{V}} = 1$ on V . For each $t \in \mathbb{R}$ holds

$$\text{var}(R1_{(a,t]}) = \text{var}(Q1_{(a,t]}) = Q1_{(a,t]}(a) - Q1_{(a,t]}(b) \leq Q1(a) = 1 .$$

2. Let $r(x, y) := - \int_a^y \frac{\partial}{\partial x} q(x, t) m(dt) = - \frac{\partial}{\partial x} Q^*1_{(a,y]}(x)$, define R with this kernel and let $s(x, y) := q(x, y) - r(x, y)$. Then $r(x, y) = s(x, y) = 0$ if $x \notin \tilde{V}$ but $y \in V$, and

$$R1_{(a,t]}(y) = \int_a^t r(x, y) m(dx) = -Q^*1_{(a,y]}(t) + Q^*1_{(a,y]}(a) = 1 - Q^*1_{(a,y]}(t)$$

by (3.29), such that $\alpha_R = \sup_t \text{var}(R1_{(a,t]}) = \sup_t \text{var}(Q1_{(a,y]}) = 1$ because of the monotonicity of $Q1_{(a,y]}(t)$ as a function of y . Furthermore,

$$R1_{\tilde{V}}(y) = 1 - Q^*1_{(a,y]}(b) = 1 \quad \text{for } y \in V \text{ by (3.29), and}$$

$$R1_W(y) = - \int_W r(x, y) m(dx) = 0 \quad \text{for } y \in V \text{ if } W \cap \tilde{V} = \emptyset.$$

In order to estimate $\text{var}_y s(x, y) = \text{var}_y(q(x, y) - r(x, y))$ we fix $\varphi \in \mathcal{T}_1(X)$ and consider

$$\begin{aligned} & \int \varphi'(y)(q(x, y) - r(x, y)) m(dy) \\ &= \int \varphi'(y) \left(q(x, y) + \int_a^y \frac{\partial q}{\partial x}(x, t) m(dt) \right) m(dy) \\ &= - \int \varphi(y) \cdot \left(\frac{\partial q}{\partial y}(x, y) + \frac{\partial q}{\partial x}(x, y) \right) m(dy) \\ &\leq C_q . \end{aligned}$$

This estimate shows that $\text{var}_y s(x, y) \leq C$ for all x . ■

The following corollary allows to apply the reasoning of part 2 of the previous one also in cases where q is not differentiable in the strict sense, but where e.g. the following condition is satisfied: There is a constant $C > 0$ such that for all $\delta > 0$ holds

$$\int |f(x + \delta, y + \delta) - f(x, y)| m(dy) \leq C \cdot \delta \quad \text{for all } x .$$

Corollary 3.21 *If Q, Q_n are (sub)-Markovian operators, $\lim_{n \rightarrow \infty} \|(Q - Q_n)h\|_1 = 0$ for all h in a dense subset of L^1 , and if $\liminf_{n \rightarrow \infty} C_{q_n} < \infty$ with C_{q_n} as in (3.31), then*

$$\text{var}(Qh) \leq \text{var}(h) + \liminf_{n \rightarrow \infty} C_{q_n} \cdot \|h\|_1.$$

Proof. $\text{var}(Qh) \leq \liminf_{n \rightarrow \infty} \text{var}(Q_n h)$ because $\lim_{n \rightarrow \infty} \|Q - Q_n\|_1 = 0$. ■

References

- [1] Baladi V., Young L.-S. On the spectra of randomly perturbed expanding maps, *Comm. Math. Phys.* 156(1993), 355-385.
- [2] Blank M.L. Small perturbations of chaotic dynamical systems. *Uspekhi Matem. Nauk.*, 44:6, 1989, p.3-28.
- [3] Blank M.L. Ergodic properties of one method of computer modeling of chaotic dynamical systems. *Matem. Zametki*, 45:4, 1989, p.3-12.
- [4] Blank M.L. Chaotic mappings and stochastic Markov chains. *Abstracts of Congress IAMP-91*, 1992, 6p.
- [5] Blank M.L. Singular phenomena in chaotic dynamical systems. *Doklady Akad. Nauk (Russia)*, 328:1(1993), 7-11.
- [6] Bunimovich L.A., Pesin Ya.G., Sinai Ya.G., Jacobson M.V. Ergodic theory of smooth dynamical systems. *Modern problems of mathematics. Fundamental trends.* v.2, 1985, P.113-231.
- [7] Doob J.L., *Stochastic processes*, Wiley, New York, 1953.
- [8] Gora P., Boyarsky A. Compactness of invariant densities for families of expanding, piecewise monotonic transformations, 41:5(1989), 855-869.
- [9] Hofbauer F., Keller G. Ergodic properties of invariant measures for piecewise monotonic transformations. *Math. Z.*, 180 (1982), 119-140.
- [10] Hunt F. A Monte Carlo approach to the approximation of invariant measures, *Random & Computational Dynamics*, 2:1(1994), 111-133.
- [11] Keller G. Stochastic stability in some chaotic dynamical systems, *Mh. Math.*, 94 (1982), 313-333.
- [12] Kifer Yu. *Random perturbations of dynamical systems*, Boston: Birkhauser, 1988.
- [13] U. Krengel, *Ergodic Theorems*, W. de Gruyter, Berlin-New York (1985).
- [14] Li T.Y. Finite approximation for the Frobenius-Perron operator. A solution to Ulam's conjecture, *J. Approx. Th.*, 17(1976), 177-186.
- [15] Lasota A., Yorke J.A. On the existence of invariant measures for piecewise monotone transformations, *Trans. Amer. Math. Soc.*, 186(1973), 481-488.
- [16] Morita T. Random iteration of one-dimensional transformations. *Osaka J. Math.* 22:3(1985), 489-518.
- [17] Pelikan S. Invariant densities for random maps of the interval. *Trans. AMS*, 281:2(1984), 813-825.
- [18] Ulam S. *Problems in modern mathematics*, Interscience Publishers, New York, 1960.